

THE BOTANICAL SOCIETY  
AND EXCHANGE CLUB  
OF THE BRITISH ISLES.

(VOL. IV. PART II).

---

REPORT FOR 1914

OF THE

BOTANICAL EXCHANGE CLUB

*(Conveniently Abbreviated for Citation REP. B.E.C.)*

BY THE

EDITOR AND DISTRIBUTOR,

R. H. CORSTORPHINE, B.Sc.

---

The Subscription, 7s 6d per annum, and Non-Contributing Members' Subscription of 5s per annum, become due on January 1, 1916, and should be sent to

G. CLARIDGE DRUCE,  
YARDLEY LODGE,  
9 CRICK ROAD, OXFORD.

Cheques for three or four years in advance would save much trouble and expense in postage.

---


Parcels for 1915 should be sent post paid, on or before 10th December 1915, to A. R. HORWOOD, Esq., MUNICIPAL MUSEUM, LEICESTER.

The Distributors and Editors for 1916 will be W. H. PEARSALL, Esq., and D. LUMB, Esq., DALTON IN FURNESS.

---

PRINTED BY T. BUNCLE & Co., ARBROATH.

November 1915.



*REPORT OF THE DISTRIBUTOR FOR 1914.*

THE total number of plants sent in for distribution shows a reduction on the two previous years. This, however, is readily accounted for by the abnormal condition of the country—many of our members having had to devote themselves to more imperative work. The number of plants sent in was 6537, contributed by thirty-four members.

The specimens were on the whole well prepared and the rules of the Club fairly well adhered to. The critical genera were all well represented, the most noticeable increase being in the genus *Erophila*. With regard to *Fumaria*, Mr Pugsley remarks "The plants were well dried, and the naming a great improvement on what obtained a few years ago." There were also a large number of good specimens of our less critical plants, which would doubtless be much valued by the newer members. Our largest contributor was Mr Robinson, whose specimens showed much care in preparation. Mr Pearsall's excellent specimens of *Hydrilla* were much appreciated.

The Rules of the Club have been reprinted, and I should like to direct the attention of the members to the remarks on the labelling of the specimens. At present the distributor receives with each packet of plants the requisite number of labels, but only *one* copy of the additional particulars and remarks of the collector regarding the plant. These remarks are generally printed in the *Report*. They are not, however, available until too late for the use of the experts who criticise the plant. This is obviously a great disadvantage, as the critic in his examination of the plant does not have before him facts as to abnormal conditions of growth or other details which might considerably influence his opinion, nor, when he is unaware of them, can he elucidate the sender's difficulties or answer his questions. Therefore, it is very necessary for those contributors who have any remarks of import to make upon their plants to send a copy of these with each label, or at least to send a sufficient number of copies to supply the critics.

The Club is greatly indebted to the following botanists for critical notes on the specimens:—Mr E. G. Baker, Mr W. Barclay, Dr Drabble, Mrs Gregory, Mr J. Groves, Mr W. P. Hiern, Rev. E. F. Linton, Dr C. E. Moss, Mr H. W. Pugsley, Rev. W. Moyle Rogers, Mr F. W. Stansfield, Dr Thellung, and Members of the Club whose names will be seen in the body of the *Report*.

R. H. CORSTORPHINE,

*Editor of Report and Distributor for 1914.*

HILLSIDE HOUSE,  
ARBROATH, NOV. 1, 1915.

## LIST OF PARCELS RECEIVED.

	No. of Specimens.
Adamson, R. S., <i>B.A.</i> ... ..	22
Bailey, Charles, <i>M.Sc.</i> , <i>F.L.S.</i> ... ..	20
Barclay, W., ... ..	34
Barton, W. C., <i>M.A.</i> ... ..	279
Bickham, S. H., <i>F.L.S.</i> , <i>J.P.</i> ... ..	98
Britton, C. E., ... ..	126
Brown, G. C., ... ..	425
Chester, G., ... ..	116
Comber, John, <i>J.P.</i> ... ..	229
Corstorphine, Mr and Mrs R. H., ... ..	292
Cryer, John, ... ..	378
Cumming, L., <i>M.A.</i> ... ..	132
Druce, G. Claridge, <i>M.A.</i> , <i>J.P.</i> ... ..	694
Groves, J., <i>F.L.S.</i> ... ..	56
Hayward, Miss Ida M., <i>F.L.S.</i> ... ..	41
Horwood, A. R., ... ..	10
Jackson, A. Bruce, ... ..	44
Johnston, Henry Halero, Col. ... ..	188
Little, J. E., <i>M.A.</i> ... ..	439
Lumb, D., ... ..	425
Marshall, Rev. E. S., <i>M.A.</i> , <i>F.L.S.</i> ... ..	190
Pearsall, W. H., ... ..	112
Riddelsdell, Rev. H. J., <i>M.A.</i> ... ..	219
Rilstone, F., ... ..	42
Robinson, F., ... ..	755
Salmon, C. E., <i>F.L.S.</i> ... ..	53
Travis, W. G., ... ..	81
Vigurs, C. C., <i>M.D.</i> ... ..	308
Waddell, Rev. C. H., ... ..	37
Waterfall, Charles, <i>F.L.S.</i> ... ..	310
Webster, Alfred, ... ..	37
Wheldon, J. A., <i>F.L.S.</i> ... ..	72
White, J. W., <i>F.L.S.</i> ... ..	238
Wilson, A., <i>F.L.S.</i> , <i>F.R.Met.S.</i> ... ..	35
Total, ... ..	6537

*Thalictrum minus* L., var. *montanum*. Dry bank, sandy soil, Tottington, v.-c. 28, July 23, 1914.—F. ROBINSON. "So I should name it."—E. S. MARSHALL.

*Thalictrum majus* Crantz. East Kennack Valley, June 19, 1914, and near Penhale, June 18 and August 12, 1914, both in the Lizard peninsula. There can be no doubt, I think, that this plant is native in the Lizard district, though it seems strange that it should be there. I think the plant is correctly named, though, as far as I know, it has never been determined by any competent authority. Is *T. majus* Crantz, of the *Lond. Cat.*, ed. x., and Druce's *List* = *T. majus* Sm. of *Bab. Man.*, ed. ix., and = *T. majus* Jacq. mentioned by Mr Salmon in last year's *Report*? The Penhale locality is a new one. Some of the plants in both places were more than four feet high. I am much indebted to Mr E. Thurston, C.I.E., for this and other Lizard plants. Coll. E. THURSTON; comm. C. C. VIGURS. "I prefer to call this *T. collinum* Wallr."—E. F. LINTON. "I have seen a similar plant on the rocky coast near Mullion. Not *T. majus*. I should refer it to *T. montanum* Wallr."—E. S. MARSHALL. "Not a *majus* form, I should say. Is it not *dunense*?"—C. E. SALMON.

*Thalictrum alpinum* L. In abundance on what is termed the "Sugar-Loaf Limestone," Cronkley Fell, Teesdale, v.-c. 65, June 9, 1914. Altitude 2000 ft.—J. CRYER.

*Myosurus minimus* L. Arable land near Madresfield, Great Malvern, v.-c. 37, May 7, 1914.—Coll. R. F. TOWNDROW; comm. C. WATERFALL.

*Ranunculus bulbosus* L. Golf Links, Askam, v.-c. 69b, June 15, 1914. I think that this could not be anything but spontaneous. I thought the plant had "flowered itself to death," but it has recovered.—D. LUMB. "This seems to be the "*Ranunculus dulcis multiplex*, Double wilde Crow-foot" figured in Johnson's *Gerard*, 957 (1633), where one reads that it "hath of late beene brought out of Lancashire unto our London gardens, by a curious gentleman in the searching forth of simples, Mr Thomas Hesketh, who found it growing wilde in the towne fields of a smal village called Hesketh, not farre from Latham in Lancashire."—C. E. SALMON. "A form with double flowers; new to me."—E. S. MARSHALL. "Good examples of the '*stare pleno*' form."—G. C. DRUCE.

*Ranunculus Flammula* L., f. *minima* Ar. Benn. Slacks in sandhills, Freshfield, v.-c. 59, July 5, 1914.—W. G. TRAVIS. "I have much the same thing from Holburn Head, Caithness. Only a *state* due to local conditions."—E. S. MARSHALL. "*R. Flammula* L."—G. C. DRUCE.

*Ranunculus sardous* Crantz. Wyke, Weymouth, v.-c. 9, July 14, 1914. Frequent round Weymouth, I believe; but this is a "new locality" for the species. The flowers are much smaller than those of *R. repens* L., but of just about the same bright polished yellow. At the Lizard, the flowers I gathered there were of a lighter lemon-yellow.—H. J. RIDDELSDELL.

*Ranunculus parviflorus* L. Hedge bank, Upper Chase Road, Malvern, v.-c. 37, May 7, 1914.—C. WATERFALL. "Yes, a very hairy form."—G. C. DRUCE.

*Ranunculus*—? [Ref. No. 2791.] Near Cowbit, Lincolnshire, July 1912.—G. C. DRUCE. "*R. Baudotii*."—J. GROVES. "Cf. *confusus*."—W. P. HIERN.

*Ranunculus heterophyllus* Weber. Pool on roadside between Great Malvern and Madresfield, v.-c. 37, May 2, 1914.—C. WATERFALL. "Yes. Carpels practically glabrous."—J. GROVES. "Is *triphyllus*."—W. P. HIERN.

*Ranunculus heterophyllus* Bab. [Ref. No. 41.] Stagnant water, Watton, v.-c. 28, May 14, 1914.—F. ROBINSON. "*R. heterophyllus* Weber. Carpels with numerous strong bristles."—J. GROVES. "Is *radians*."—W. P. HIERN.

*Ranunculus heterophyllus* Weber, var. *triphyllus* Wallr. Portbury marshes, N. Somerset, May 29, 1914.—J. W. WHITE. "Is *radians*."—W. P. HIERN. "*R. heterophyllus*. The carpels are bristly, whereas those of *R. triphyllus* Wallr. are described as 'glaberrimis nitidis.'"—J. GROVES.

*Ranunculus heterophyllus* Weber, var. *submersus* Hiern. Marsh ditch, Portbury, N. Somerset, June 3, 1902.—J. W. WHITE. "Is *trichophyllus*."—W. P. HIERN. "I am doubtful about this. It has more rigid leaves than is usual in *R. heterophyllus submersus*. The entirely undeveloped heads of carpels suggest hybridity. I do not think we can properly cite Hiern as the authority for var. *submersus* of *R. heterophyllus*, as in his paper on the group he treated the whole genus *Batrachium* as one species. Moreover, his form No. 30 *submersus* was not ranged under the head of *R. heterophyllus*."—J. GROVES.

*Ranunculus peltatus* Schrank, var. *truncatus* (Hiern). Birkdault, v.-c. 69b, June 13, 1914. This grows in the same ditch as the form distributed last year [Ref. No. 380], and seems intermediate between it and var. *truncatus*.—W. H. PEARSALL. "*R. truncatus*."—W. P. HIERN. "Yes, I suppose this curious plant must be so labelled, but I have never seen *truncatus* with similar leaves."—J. GROVES.

*Ranunculus pseudo-fluitans* Bab. In the River Ribble, near Preston, South and West Lancashire, v.-c. 59 and 60, July 1914. This is probably the plant recorded from about ten miles higher up the river in the *Flora of Stonyhurst* as *R. fluitans* Lam., "not flowering" (see *Flora of West Lancashire*, p. 129). During this last dry summer flowers were freely produced near Preston, the river being exceptionally low.—A. WILSON. "My specimen is too meagre for determination; fruit and floating leaves absent, and only one flower."—E. S. MARSHALL. "From this specimen, I should have thought a form of *R. fluitans*, but I should like to see more of it."—J. GROVES. "Cf. *R. Bachii*."—W. P. HIERN. "This appears to be identical with the Batrachian *Ranunculus* sent to the Club in 1904 by the same collector from the River Wharfe, W. Yorks, under the name of *R. pseudo-fluitans* Hiern. The latter plant was considered by Messrs Groves to be a small river form of the *peltatus* group. Examples of both gatherings agree with descriptions of *R. pseudo-fluitans* except in the characters of robustness and the average length of the peduncles, and both curiously agree in the fact of being completely sterile! If this latter characteristic is not unusual, it would probably be as well to publish a description of this form as a new species."—C. E. BRITTON.

*Ranunculus Baudotii* Godr.? Brackish ditch inside sea wall between Sidlesham and Pagham, W. Sussex, v.-c. 13, June 15, 1914.—J. E. LITTLE. "*R. Baudotii*."—C. E. BRITTON, J. GROVES, and W. P. HIERN.

*Ranunculus Baudotii* Godr. Roose, v.-c. 69b, May 22, 1914. Very plentiful in brackish water near the coast.—W. H. PEARSALL. "A small form of this, I think."—E. S. MARSHALL. "Yes."—J. GROVES. "Cf. *confusus*."—W. P. HIERN.

*Aquilegia vulgaris* L. Marsh, Carbrooke Fen, v.-c. 28, May 19, 1914.—F. ROBINSON.

*Papaver hybridum* L. Near High Down, Hitchin, Herts, v.-c. 20, August 1, 1914.—J. E. LITTLE.

*Corydalis lutea* DC. Newquay, Cardiganshire, May 1914.—J. W. WHITE.

*Corydalis claviculata* DC. Clophill, Beds, v.-c., 30, April 25, 1914. Not recorded by Abbot (*Flora Bedfordshire*, 1798).—J. E. LITTLE. "Small specimens. It is a plant very responsive to conditions of shade or exposure. The var. *minor* R. & F. *Flore de France*, p. 188, from schistose soil, is only 4 to 12 cm. high. I have seen such on recently burnt commons. I found it at Woburn in 1874."—G. C. DRUCE.

*Fumaria paradoxa* Pugsley. Orig. near Reigate Hill, Surrey, 1912, Hort. Reigate, 1914. It is difficult to imagine a more beautiful Fumitory, when in a fresh state, than the one now distributed. It is a mystery how the plant reached the cultivated ground near the farm on Reigate Hill where I gathered it on June 16, 1912. Not far away were *Thlaspi arvense*, *Lotium temulentum*, and two or three plants of *Hyoscyamus*. Could the *Fumaria* have been accidentally introduced with potatoes from Cornwall?—C. E. SALMON. "Yes; I have rarely seen cultivated specimens of *Fumaria* so satisfactory as this."—H. W. PUGSLEY.

*Fumaria capreolata* L., var. *Babingtonii* Pugsley. Blackhead, Co. Clare, June 1909.—G. C. DRUCE. "Showing no fruit, but no doubt correctly named."—H. W. PUGSLEY.

*Fumaria Boraei* Jord. [Ref. No. 1168.] Chobham, Surrey. May 31, 1914.—C. E. BRITTON. "Yes; under type, I believe."—E. S. MARSHALL. "Correct; near Jordan's type, but with rather small flowers and fruits. This is a rare plant in Surrey."—H. W. PUGSLEY.

*Fumaria* —? Odiham, Hampshire, July 1903. (See *Report* 1903, p. 9.)—Coll. C. E. PALMER; comm. G. C. DRUCE.

*Fumaria Bastardi* Bor. (*F. confusa* Jord.). Arable land near Llangollen, Denbigh, v.-c. 50, September 20, 1914.—C. WATERFALL. "Yes. The sheet sent is var. *hibernica* Pugsley. Easily distinguishable by its dark-tipped upper petal."—H. W. PUGSLEY.

*Fumaria officinalis* L., var. ? This appeared in great abundance in a ploughed field at Charlestown, Baildon, v.-c. 64, June 1, 1914.—J. CRYER. "A narrow-leaved form; but I do not see the varietal character."—E. S. MARSHALL. "The sheet sent is a floriferous, early flowering form of *F. officinalis* L., var. *Wirtgeni* Haussk., but some of the fruits enclosed in the envelope are different from the majority, and appear to have come from a plant of *F. officinalis* type."—H. W. PUGSLEY.

*Barbarea verna* Ascher. Garden weed, Colchester, v.-c. 19, May 23, 1914.—G. C. BROWN. "Rightly named."—C. E. SALMON. "Yes; very characteristic."—A. B. JACKSON.

*Barbarea verna* Ascher. Railway, Askam, v.-c. 69b, July 5, 1914. Woods gives as a character of this plant "auricles ciliate." Is any member able to say whether this is constant and diagnostic? There were what seemed to me two plants growing together, and I feel certain that I have failed to separate them; in all probability I have paid too much attention to "ciliate auricles."—D. LUMB. "No; this

I should call *B. intermedia* Bor."—C. E. SALMON. "Looks right."—E. S. MARSHALL.

*Barbarea vulgaris* Bor. Railway, Askham, v.-c. 69b, July 5, 1914. I have named this with some doubt. All these plants may possibly belong to the other set.—D. LUMB. "*B. intermedia* Bor., I think, but specimens badly dried."—A. B. JACKSON. "Is *B. intermedia* Bor."—A. THELLUNG.

*Barbarea vulgaris* R. Br., var. *campestris* Fr. Nov. Fl. Suec., p. 205, 1828. Stiff clay soil on railway embankment between Alperton and Sudbury, Middlesex, May 22, 1914. This, the commonest variety of *B. vulgaris* in Britain, is distinguished by its slightly spreading or secund pods, but it passes gradually into the var. *arcuata* Fr. (pods arcuate) on the one hand, and into var. *sylvestris* (pods adpressed) on the other.—A. B. JACKSON.

*Arabis hirsuta* Scop., var. Symond's Yat, W. Glos, May 28, 1913. Of course only a slight variety, with the pods somewhat spreading. As a rule, the species has closely adpressed pods, even in shady woods; the variety is not due, as it seems, to the presence of shade. There is a parallel variety in *Barbarea vulgaris*. The form occurs also on the Great Doward, v.-c. 36. The sparseness of the hairs is probably due to the less exposed conditions.—H. J. RIDDELSDELL. "I collected this (same place and date) and thought it untypical. No special name suggested so far."—E. S. MARSHALL. "This seems nearest to var. *gracilescens* (Jord.) R. & F. Fl. Fr. i., 216. Plante assez élevée (3—5 dm.) mais à tiges grêles flexueuses; feuilles d'un vert clair, les caulinaires tronquées ou légèrement subcordées, lancéolées ou oblongues-lancéolées, acutiuscules, nombreuses, à 4 dents souvent saillantes; siliques (25—35 mm.) étroitement linéaires, disposées en grappe allongée, lâche. I have the same form from Wells, Somerset; Culford, Suffolk."—G. C. DRUCE.

*Arabis scabra* All. Clifton, W. Glos., v.-c. 34, April 27, 1914.—W. C. BARTON.

*Arabis alpina* L. North side of the Cuchullins, Skye, June 1910.—G. C. DRUCE and T. H. LEACH. This was from a different locality on the Cuchullins to that which was discovered by Mr H. Hart in 1887, and is, I believe, the second time it has been gathered in the British Isles. Mr Hart's specimens, gathered on his wedding tour in the first week in July, are in fruit; ours gathered in June are in good flower. The plant is very local and requires climbing to reach (2700—2800 ft. alt.), growing on damp rock ledges. Mr A. H. Evans and Mr T. H. Leach (my godson) were with me, and the latter was the first to spot the prize, for we were systematically working the

corrie in sections. We did not see it on Scur Alister, where it is believed Mr Hart originally found it.—G. C. DRUCE.

*Arabis petraea* Lam., var. *hispida* DC. Ben Hope, W. Sutherland, July 1907. This hispid variety of *A. petraea* from Ben Hope differs from the plant of the Cairngorms and Snowdon in having much larger flowers, in this point resembling my var. *grandifolia* from Ben Laoigh; in fact, a few plants referable to that variety were found there. Mr Arthur Bennett referred my *grandifolia* to *A. petraea*, var. *ambigua* Fries Mantissa iii., 77; the vague definition "elatior, foliis radicalibus lyrato-sinuatis caulinis subdentatis radice tenuiori" does not give the essential characters of the Ben Laoigh plant I designated var. *grandifolia*, which must stand for the Ben Laoigh plant. The var. *ambigua* Fries, *A. ambigua* DC. *Syst.* i., 231, is chiefly Siberian and Unalaskan and is not a perennial, and he makes no mention of size of leaves or flowers.—G. C. DRUCE.

*Arabis glabra* Bernh. (*perfoliata* Lam.). Dry heath amongst gorse, Barnham Common, v.-c. 28, August 22, 1914.—F. ROBINSON.

*Cardamine impatiens* L. Seedlings. Occurs as a weed on several gardens in an allotment near Kettering, v.-c. 32, Jan. 17, 1914.—G. CHESTER.

*Erophila*. [Ref. No. 2041]. Fairford, Glos., April 1904. This is a not uncommon plant of our oolitic areas in Oxfordshire and Gloucestershire, and is, I think, *E. Ozanoni* Jord. *Diagn.* 231, et *Icones*, t. 5, n. 17.—G. C. DRUCE. "Some specimens rather closely approach *E. praecox* DC."—E. S. MARSHALL.

*Erophila*—? [Ref. No. 50]. Edge of salt marsh near Montrose, v.-c. 90, May 16, 1914.—R. and M. CORSTORPHINE. "This is a form of *E. majuscula* Jord., and apparently is the *Draba majuscula* R. & F., var. *occidentalis* R. & F. = *E. occidentalis* Jord."—C. E. BRITTON. "Depauperate *E. stenocarpa* I strongly suspect."—E. S. MARSHALL.

*Erophila*—? [Ref. No. 56]. Roadside near Restennet, v.-c. 90, May 2, 1914.—R. & M. CORSTORPHINE. "I think there are two plants here. The bulk is nearly glabrous, the few hairs mostly simple. The silicle measurements and narrow leaves suggest placing it under *Draba glabrescens*, var. *erratica* Rouy et Fouc."—J. A. WHELDON. "A peculiar little plant. Pods much reticulate, veined as they mature. Leaf surface nearly glabrous, except near the margins."—E. S. MARSHALL.

*Erophila verna* E. Meyer. [Ref. No. 4]. Wall under trees, in a dip in the road from Hook Norton to Wigginton, Oxon, April 14,

1914. This bushy, much branched *Erophila* occupied a section of the wall top, pretty much to the exclusion of everything else. Perhaps it should come under *E. verna* agg.—H. J. RIDDELSDELL. "This seems to occupy an intermediate position between *E. præcox* and *E. majuscula*, though considerably nearer the latter. I would name it *E. obovata* Jord."—C. E. BRITTON. "Petals large; pods broad, round-topped. Resembles what I have as British *E. majuscula* Jord."—E. S. MARSHALL. "These come under *E. præcox* DC."—G. C. DRUCE.

*Erophila verna* E. Meyer, var. [Ref. No. 5]. Wall, fully exposed, between Hook Norton and Wigginton, Oxon, April 14, 1914. A plant with darker foliage than Ref. No. 4, narrower pods, and quite distinct habit.—H. J. RIDDELSDELL. "I think this must be referred to *E. majuscula* Jord., from which it differs chiefly by the narrower leaves. It is probably the var. *occidentalis* R. & F. = *E. occidentalis* Jord."—C. E. BRITTON.

*Erophila verna* Meyer: [Ref. No. 54]. Field track near Rescobie, v.-c. 90, May 2, 1914.—R. & M. CORSTORPHINE. "Small plants, which seem to me to be nearer *E. stenocarpa* Jord."—E. S. MARSHALL.

*Erophila majuscula* Jord. [Ref. No. 52]. The Lurgies near Montrose, v.-c. 90, April 27, 1914.—R. & M. CORSTORPHINE. "*E. majuscula* Jord."—J. A. WHELDON. "Yes."—C. E. SALMON. "Correctly named."—C. E. BRITTON. "*E. stenocarpa* Jord., mostly very typical."—E. S. MARSHALL.

*Erophila majuscula* Jord. [Ref. No. 52b]. Near Bridge of Dun, v.-c. 90, May 14, 1914.—R. & M. CORSTORPHINE. "Yes, I suppose best so labelled, but silicles less rounded above than usual, giving more the shape of those of *E. stenocarpa*. But the large petals and width of silicles quite exclude that."—J. A. WHELDON. "Correct, I believe."—C. E. SALMON. "Correctly named."—C. E. BRITTON. "No, my specimen is good *E. stenocarpa* Jord."—E. S. MARSHALL.

*Erophila majuscula* Jord. ? [Ref. No. 83]. Cultivated ground on Ash Brook, St Ippolyts, Herts, v.-c. 20, April 10, 1914. The most luxuriant examples grow to a considerable size (14 cm), and have large rosettes of broad leaves.—J. E. LITTLE. "I do not recollect seeing anything quite like this. Conf. *E. affinis* Jord. (*Draba leptophylla*, var. *australis* R. & F.). Hairs bifid (rarely trifid or simple); flowers large; silicle oblong; calyx and petals often suffused with violet-rose."—J. A. WHELDON. "*E. occidentalis* Jord."—C. E. BRITTON. "Foliage curious. Near *E. verna*; but I am not able to name this."—E. S. MARSHALL.

*Erophila verna* E. Meyer, d. *majuscula* (Jord.). Wall top and edge of footpath, the Marine Drive, The Great Orme, v.-c. 49, March 12, 1914.—C. WATERFALL.

*Erophila stenocarpa* Jord. Field at St Ippolyts, near 2nd milestone for Hitchin, Herts, v.-c. 20, April 24, 1914. Cornfield, about a mile away from Ref. No. 60 (*Report* 1913, p. 450), in similar stony loam. As with Ref. No. 60, there was an admixture of plants with less characteristic silicles.—J. E. LITTLE. "Right, I think."—C. E. SALMON. "One specimen is certainly right (rather small); the others come between that and *E. verna* (*vulgaris* DC.)."—E. S. MARSHALL.

*Erophila stenocarpa* Jord. Sandy cultivated ground, Mauldon, Beds, v.-c. 30, April 25, 1914. The later stage of a plant similar to the preceding, and, like it, mixed with less characteristic forms.—J. E. LITTLE. "Yes, I think so, but not an extreme example." J. A. WHELDON. "Yes."—E. S. MARSHALL.

*Erophila verna* E. Meyer, var. *stenocarpa* (Jord.). [Ref. No. 6.] From another part of the wall on which No. 4 occurred, between Hook Norton and Wigginton, Oxon, April 14, 1914. Some of this small gathering seems to fit *stenocarpa* very well, but some of it looked as if it had a touch of No. 4 in it.—H. J. RIDDELSDELL. "Four of my five specimens are *E. stenocarpa* Jord.; the other has shorter pods and approaches *E. verna*."—E. S. MARSHALL.

*Erophila stenocarpa* Jord.? [Ref. No. 53]. Field side near Lunanhead, v.-c. 90, May 2, 1914.—R. & M. CORSTORPHINE. "Silicles  $5 \times 2$  mm. In *E. stenocarpa* they frequently attain 7 mm. in length, while only  $1\frac{1}{2}$ — $1\frac{3}{4}$  mm. broad. I would suggest that this is a form of *E. brevipila* Jord."—J. A. WHELDON. "I think so; a weak small podded state."—E. S. MARSHALL.

*Erophila obovata* Jord.? [Ref. No. 55]. Wall top near Clocksbriggs, v.-c. 90, May 2, 1914.—R. & M. CORSTORPHINE. "I do not know that. Leaves remarkably glabrous at this stage."—E. S. MARSHALL.

*Erophila præcox* DC. [Ref. No. 3]. Wigginton, Oxon, April 11, 1914. The wall tops of this neighbourhood are covered with *Erophila*, a fair proportion of which is *E. præcox*. These elusive micro-species are all the more difficult to determine, in a great number of individual cases (some of the specimens now sent are only doubtfully *præcox*), because the colonies are by no means homogeneous; and there is much obvious transition between species, to say nothing of highly probable crossing. Another difficulty arises from the fact that pods vary in shape even on a single plant. The best and unmistakable *præcox* runs

small.—H. J. RIDDELSDELL. "Yes, mostly a tall drawn up form."—E. S. MARSHALL.

*Erophila præcox* DC.? Sandhills, Askham, v.-c. 69b, April 12, 1914. This seems to be the same plant as I sent last year.—D. LUMB. "Clearly *E. præcox* DC."—C. E. BRITTON. "Yes, the usual small coast form."—E. S. MARSHALL. "Yes, a new county record for 69b."—G. C. DRUCE.

*Erophila præcox* DC. [Ref No. 58]. Sands of Barry, v.-c. 90, April 26, 1914.—R. & M. CORSTORPHINE. "Yes, the normal plant of coast sands."—E. S. MARSHALL. "Yes, for one specimen at least. A new county record."—G. C. DRUCE.

*Erophila spathulæfolia* Jord.? (fide J. A. Wheldon). Gravel pit, St Ippolyts, Hitchin, Herts, v.-c. 20, April 4, 1913, and March 17, 1914. I sent this plant to the *Watson B.E.C.* 1913-14, named by Mr J. A. Wheldon as *E. spathulæfolia* Jord. I think it is reasonably certain that it belongs to Rouy and Foucaud's sub-sp. vii. *Draba glabrescens* Rouy and Fouc. (*Erophila glabrescens* Jord.); but I have not felt able to carry the limitation further. The specimens now sent are the result of selection three times repeated, ending in the elimination of a very much larger number of plants with more numerous bifid hairs. I have not succeeded in finding any other locality in this neighbourhood for this form with long sparse mostly simple hairs.—J. E. LITTLE. "The examples sent under this name belong to that section of aggregate *E. verna* distinguished by the hairs being predominantly simple rather than branched. These specimens come very close to Jordan's description of *E. spathulæfolia*, but differ in the glabrous scapes, shorter pedicels, and narrower silicules. For these reasons I do not think these plants can be referred to *E. spathulæfolia* Jord."—C. E. BRITTON. "I do not know that. The leaves are certainly remarkable."—E. S. MARSHALL.

*Cochlearia groenlandica* L. Coast at Boddin, Forfar, v.-c. 90, June 3, 1914. N.C.R. for Forfar.—R. & M. CORSTORPHINE. "Certainly *C. groenlandica*."—E. S. MARSHALL. "Yes, a new county record, I believe."—G. C. DRUCE.

*Sisymbrium Sophia* L. Sainfoin field, Thetford, v.-c. 28, June 1, 1914.—F. ROBINSON.

*Sisymbrium pannonicum* Jacq. = *S. altissimum* L. Waste heap, North of Welwyn Tunnel, Herts, v.-c. 20, June 16, 1913. I send as a record. The waste heap is now cultivated.—J. E. LITTLE. "Yes, the older name is *S. altissimum* L., a frequent alien."—G. C. DRUCE.

*Sisymbrium orientale* L. (= *S. Columnae* Jacq.). Par Harbour, East Cornwall, June 19, 1914. I send this common alien because the plants are mostly young ones, and show the lower foliage better than I have ever seen before.—C. C. VIGURS. "Yes."—G. C. DRUCE. "The var. *subhastatum* (Willd.) Thell."—A. THELLUNG.

*Brassica elongata* Ehrh. Roadside, Billingshurst, Sussex, July 22, 1914.—A. WEBSTER. "The sub-sp. *persica* (B. & R.) Thell."—A. THELLUNG.

*Brassica Erucastrum* Vill. [Ref. No. 40]. Sandhills by sea, Yarmouth, N. Devon, v.-c. 27, May 10, 1914.—F. ROBINSON. "Is *Sisymbrium orientale* L."—A. THELLUNG.

*Brassica balearica* Loisel. ? Railway side, Askham, v.-c. 69b, July 2, 1914.—D. LUMB. "*Brassica juncea* (L.) Coss."—A. THELLUNG.

*Diplotaxis tenuifolia* DC. Railway cutting, Thetford, v.-c. 28, June 15, 1914.—F. ROBINSON. "This must, I think, be referred to *D. muralis*, var. *Babingtonii* (Syme). In *tenuifolia* the flowers are distinctly stalked at the time of full flowering, and the pods are more distinctly narrowed at both ends. This is the biennial form."—G. C. DRUCE.

*Diplotaxis*—? [Ref. No. 716]. On cotton seed refuse, Hythe Quay, Colchester, v.-c. 19, June 1914.—G. C. BROWN. "*Erucaria myagroides* Halac."—G. C. DRUCE.

*Bursa pastoris* Weber, var. *bifida* Druce. Hort., Oxon, July 1908. This plant, which approaches *macrocarpa* in having a few of the silicules with curved sides, has persisted for the last 20 years as a weed in the Botanic Garden, Oxford.—G. C. DRUCE.

*Bursa pastoris* Weber, var. *bifida* Druce. Seed from plants from the Botanic Garden, Oxford (1890), Hort. Druce, 1913. The deep sinus and shape of capsule and leaves remain practically unaltered in culture.—G. C. DRUCE. "This is the form of *Capsella Bursa-pastoris* that F. M. Mott called var. *bifida*. It is one of the best marked varieties of Shepherd's Purse."—C. E. BRITTON.

*Lepidium latifolium* L. River Lavant, near Appledram, W. Sussex, v.-c. 13, Sept. 23, 1913.—J. E. LITTLE.

*Lepidium Smithii* Hook., var. *alastostyla* Towns. Garden specimen from a 2-year-old plant raised from seed collected at Redbridge, near Southampton, Sept. 1901. Hab.—rough hedge banks on the coast. See *Report* 1903, p. 10.—Coll. FREDK. TOWNSEND; comm. G. C. DRUCE.

*Lepidium neglectum* Thell. Waste ground, Askam, v.-c. 69b, August 9, 1914. I name these plants thus with some doubt.—D. LUMB. "Yes."—C. E. SALMON and A. THELLUNG.

*Thlaspi alpestre* L., var. *occitanum* (Jord.). Roadside bank, Llanrwst, Carnarvon, v.-c. 49, April 1878.—J. COMBER. "This is not *Thlaspi occitanicum* Jord. The specimens are not sufficiently developed to show the characters of the capsules, but judging by the evidence afforded by the habit, foliage, and flowers, this is *Thlaspi virens* Jord.—C. E. BRITTON. "Wrongly named as Jordan's plant (*T. occitanicum*). I know it well in the Llanrwst district but am not sure if it differs from type."—E. S. MARSHALL.

*Teesdalia nudicaulis* Br., var. *minor*. Dry heath by Ringmere, Roudham, v.-c. 28, April 26, 1914. This tiny plant grows in the thick moss on Roudham Heath very sparsely, nothing but the tiny head of flowers shows above the moss. It is confined as far as I see to the dry bank of one of the meres which formed part of the old Fen sea.—F. ROBINSON. "Nothing varietal about this. Small states are frequent on poor soil."—E. S. MARSHALL. "Not a variety, only a condition."—G. C. DRUCE.

*Hutchinsia petraea* Br. Sparingly scattered over several boulders near Lovers' Leap, Dovedale, Derbyshire, v.-c. 57, April 14, 1914.—G. CHESTER. Also sent from limestone rocks, The Great Orme, Carnarvon, v.-c. 49, March 11, 1914.—C. WATERFALL.

*Reseda lutea* L., var. *pulchella* J. Muell. [Ref. No. 1375]. Worms Heath, Surrey, Sept. 6, 1914. This is a slender more refined form of *R. lutea*, abundantly branched, with leaf-segments elongated, flat, linear, minutely callous toothed, racemes narrow and flowers smaller than type. My plant agrees well with the original description of this var. by J. Mueller in his monograph of the *Resedaceae*, and also accords well with the figure in Reichenbach's *Icones* where it is published under the name of *R. gracilis* Tenore. There is no material quite like my plant in the collections at South Kensington.—C. E. BRITTON. "The capsules on my specimens are too young to see if they are papillose; if smooth the plant may be var. *Lecoqii* J. Muell. *R. gracilis* of Reichenbach's *Icones* is not identical with *R. gracilis* Tenore, which was recorded from Wandsworth by A. Irvine in *Fl. Surrey*."—G. C. DRUCE.

*Reseda lutea* L., var. *stricta* (Persoon, as a species). Wytham, Berks, Sept. 1906. I do not think this deserves more than varietal rank. It still persists at Wytham Mill, Berks.—G. C. DRUCE.

*Helianthemum Chamæcistus* × *polifolium*. [Ref. No. 3350] Root from Purn Hill, Bleadon, N. Somerset, v.-c. 6, where it grows mixed

with both species. Flower garden, West Monkton, May 27, 1914. A very pretty plant.—E. S. MARSHALL.

*Helianthemum canum* Baumg., var. *vineale* (Pers.). On the Sugar Limestone of Cronkley Fell, Teesdale, v.-c. 65, June 9, 1914. Altitude 2000 feet. Too early for flower.—J. CRYER.

*Viola montana* L. Wood Walton Fen, Hunts, July 1908. All these were named *montana* by Mrs Gregory in that year. I am not quite sure whether she would not now refer them to *V. stagnina* × *canina*. *V. montana* L. itself is not more than a hybrid. With these specimens were others more closely approaching *canina* and *stagnina*, and again others which appeared to be ternary hybrids.—G. C. DRUCE.

*Viola sylvestris* Kit. [Ref. No. 25]. Hedge banks, Redbill, Walton, v.-c. 28, April 19, 1914.—F. ROBINSON. "*Viola sylvestris* Lam., emend. Reichb., var. *punctata* Druce. Note the thick, furrowed spur; also the central shoot lengthening and flowering."—E. S. GREGORY.

*Viola sylvestris* Lam., emend. Reichb. Roudsea Wood, N. Lancs, April 30, 1914. Rare in N. Lancs, where the var. *punctata* is quite common.—W. H. PEARSALL. "Yes, typical *V. sylvestris*, and the most representative gathering I've seen for many years."—E. S. GREGORY.

*Viola sylvestris* Kit., var. *punctata* Druce. Sturts' Copse, Oxon, March 1913. A common form of our calcareous woods.—G. C. DRUCE.

*Viola Riviniana* Reichb. Hitch Wood, Herts, v.-c. 20, April 13, 1912. Det. E. S. GREGORY.—J. E. LITTLE.

*Viola Riviniana* Reichb., forma *minor*. Kirkby Moor, v.-c. 69b, May 16, 1914. This was named "floriferous forma *minor*" by Mrs Gregory in 1913.—D. LUMB.

*Viola Riviniana* Reichb., forma *nemorosa* Neuman. Hitch Wood, Herts, v.-c. 20, May 13, 1914. Mrs Gregory *in lit.* (June 21, 1914) writes:—"I congratulate you on having collected the best set of *Viola Riviniana*, forma *nemorosa*, that I have ever seen. It is strange how near some of the plants are to *V. silvestris* (type)."—J. E. LITTLE.

*Viola Riviniana* Reichb., var. *diversa* Greg. On the banks of the Calder, Wakefield, v.-c. 63, May 12, 1914. Mrs Gregory says:—"There is little doubt that they represent my var. *diversa* of *V. Riviniana*."—J. CRYER.

*Viola canina* L. [Ref. No. 37.] Sandy heath land by sea, North Denes, Yarmouth, v.-c. 27, May 9, 1914.—F. ROBINSON. "Yes, the variety *pusilla* Bab."—E. S. GREGORY.

*Viola canina* L., var. *lanceolata* Martin-Donos. Lake Windermere (shingly stones), N. Lancs., May 16, 1914. This is locally abundant, but I can find it only in such situations, either growing among grass on the stones or invading them on its own account.—W. H. PEARSALL. "Not at all like the luxuriant plants from Woodwalton Fen, Hunts, so named for me (on the spot) by Mrs Gregory. Perhaps the large-flowered var. *macrantha* Gren. & Godr., from Berrow, N. Somerset, which seems to me the same as the usual plant of the Scottish Highlands."—E. S. MARSHALL. "Yes, very near akin to the Norfolk plant described on page 82 of *British Violets*."—E. S. GREGORY. "Yes, Mr Pearsall showed it to me *in situ*, and it approaches closely our Oxon plant."—G. C. DRUCE.

*Viola canina* L.  $\times$  *V. lactea* Sm. Open downs near the sea, St Agnes, Scorrier, v.-c. 1; (a) with flowers, May 11; (b) with fruit, July 9, 1914. So named by Mrs Gregory from fresh material forwarded by Mr Druce in May. The commonest form of violet on the downs from which specimens were taken, the only others noticed being *V. lactea* Sm. (in small quantity) and a few plants of *V. Riviniana* Reichb.—F. RILSTONE. "I very much doubt this identification, and would rather name it *V. lactea*  $\times$  *Riviniana*."—E. S. MARSHALL.

*Viola lactea*  $\times$  *Riviniana*. [Ref. No. 3535]. Root from Crowcombe, Heathfield, S. Somerset, v.-c. 5; flower garden, West Monkton, May 18, 1914. Like the Tidenham Chase (W. Glos.) plant, this flowers very freely, but never fruits. A good intermediate between the parents, with which it was found.—E. S. MARSHALL. "Yes."—E. S. GREGORY.

*Viola hirta* L. Open wood on sandy soil, South Pickenham, v.-c. 28, April 23, 1914.—F. ROBINSON. "*V. hirta* var. *hirsuta*, in the semi-cleistogamous stage. The plants received by me appear to belong to the new form (*luteo-canescens*) of this variety, lately so named by Dr Moss and myself."—E. S. GREGORY.

*Viola hirta* L. These specimens, with very large and conspicuous flowers, were found growing on the roadside about two miles from Carnforth going towards Silverdale, v.-c. 60, April 11, 1914. In many of the flowers the hook of the spur was scarcely indicated. Mrs Gregory says:—"The *V. hirta* you send reminds me of one which I found last year near Torquay. The shape and size of the flowers suggest a giant race."—J. CRYER. "Yes, petals unusually broad and rounded I think."—E. S. MARSHALL.

*Viola hirta* × *odorata*. [Ref. No. 715]. Clayey hedgerow, Edwardstone, W. Suffolk, v.-c. 26, April 16, 1914. A few more sheets of this hybrid from a station about three-quarters of a mile from my Ref. No. 81, sent to Club last year. Rather stunted owing to exposure from hedge-cutting.—G. C. BROWN.

× *Viola multicaulis* Jord. = *V. hirta* × *odorata*. [Ref. No. 13]. Bladon, Oxon, with both parents, April 1914. Some plants were of a very large size.—G. C. DRUCE. "*Viola hirta* × *odorata* = × *multicaulis*."—E. S. GREGORY.

*Viola*—? [Ref. No. 333]. Garden weed, Dalton, v.c. 69b, August 13, 1914. These were growing entangled among the *arvatica* and a pale-flowered cultivated pansy. They may be very luxuriant *arvatica*, but the large flowers seem to point to another species or to a possible smirch with the garden pansy.—D. LUMB. "The broad leaved plants in the set are yellow flowered *Lloydii*; the narrow leaved ones approach *Provostii* Boreau. These narrow leaved plants are possibly hybrids, but there is not any satisfactory evidence for this."—E. DRABBLE.

*Viola Deseglisei* Jord. [Ref. No. 67]. Near Brechin, v.-c. 90, August 16, 1914.—R. & M. CORSTORPHINE. "Yes. Very beautiful examples. The habit as shown in the longer spreading specimens is quite characteristic. In some cases the leaves and stipules are less toothed than usual."—E. DRABBLE.

*Viola Deseglisei* Jord. [Ref. No. 659]. Field, Layer Marney, N. Essex, v.-c. 19, June 7, 1914. Apparently coming under this, though the leaves are rather broad. The branching habit and thin light green leaves would seem typical. I suppose, if correct, it would be a new county record for v.-c. 19.—G. C. BROWN. "I should say *V. arvensis* Murr., var. *ruralis* Corb."—J. CRYER. "Some of these—the narrow leaved plants—are certainly *Deseglisei*; the broad leaved specimens are probably *Deseglisei* also, but approach *ruralis* somewhat."—E. DRABBLE.

*Viola obtusifolia* Jord. (small form). [Ref. No. 660]. Field, Layer Marney, N. Essex, v.-c. 19, June 7, 1914. A new county record for v.-c. 19, if correct.—G. C. BROWN. "No, this is *ruralis*."—E. DRABBLE.

*Viola obtusifolia* Jord. [Ref. No. 1198]. Send, Surrey, June 21, 1914.—C. E. BRITTON. "Yes, *obtusifolia* Jord."—E. DRABBLE.

*Viola obtusifolia* Jord. [Ref. No. 661]. Field, Baydon, E. Suffolk, June 1, 1914. A new county record for v.-c. 25, if correct.

G. C. BROWN. "Yes, *obtusifolia*, but curiously stout-stemmed, coarse-looking plants."—E. DRABBLE.

*Viola arvensis* Jord. Garden weed, Dalton, v.-c. 69b, August 13, 1914. These were growing in very stiff, well-manured soil. They were very luxuriant and a tangled mass of stems.—D. LUMB. "The smaller plants with widely spreading peduncles are *arvensis*; and as there seem to be all gradations between these and the larger plants, the latter must be *arvensis*, too, I suppose. But the large plants are much bigger leaved and less zig-zag than any I have previously passed as *arvensis*."—E. DRABBLE.

*Viola lutea* Huds., with its form or variety *amoena* (Symons). Nos. 1 and 2 on each sheet were gathered at Malham, v.-c. 64, where they grow together. No. 3 was gathered in Teesdale, v.-c. 65, where it is very plentiful and the prevailing form. June 1914.—J. CRYER. "Yes, very beautiful specimens. It is interesting to notice the much broader leaves and petals in these plants than in Mr Waterfall's Cumberland specimens. The large Teesdale plant seems to be Mr E. G. Baker's sub-var. *insignis*."—E. DRABBLE.

*Viola lutea* Huds, b. *amoena* (T. F. Forst.). Moorlands, ascent of Catterpallot, near Melmerby, v.-c. 70, Aug. 3, 1914.—C. WATERFALL. "Yes, *amoena* is simply a blue flowered *lutea*. These are very good specimens, but there is usually an extensive underground development of the branches as in *lutea*."—E. DRABBLE.

*Polygala vulgaris* L., var. *Ballii* Ostenfeld (Not *P. Ballii* Nyman nomen). Bettyhill, Sutherland, July 1907. Practically identical with Faroese plants sent me by Dr Ostenfeld. He thought these were *P. vulgaris*, var. *grandiflora* Bab. = *P. Babingtonii* Druce, but they differ materially from the Ben Bulbin plant. Nyman, under *P. vulgaris* as a synonym (sine descriptione), gave *P. Ballii*, evidently meaning by this name the Irish plant. Therefore, if we use the name it must be as var. *Ballii* Ostenfeld, i.e., not of Nyman. I found it locally on the Limestone at Ardrahan, Galway, and Inchnadamph, Sutherland W. in 1907. Ostenfeld says it is common in the Faroes.—G. C. DRUCE. "I have carefully compared my specimen with examples from Wye Downs, Kent, and also with specimens from the hills around Grassington, Yorkshire, and I can find no characters which separate it from type *Polygala vulgaris* L."—J. CRYER.

*Dianthus prolifer* L. Shingle, Pagham, W. Sussex, v.-c. 13, June 13, 1914. See Arnold's *Sussex Flora* 1907, p. 16.—J. E. LITTLE.

*Silene latifolia* Rendle & Britten, var. *hirsuta* Gray. [Ref. No. 642]. Cultivated ground, Whatfield, W. Suffolk, v.-c. 26, June 1,

1914. Apparently identical with the plant distributed by Mr M<sup>c</sup>T. Cowan, jun., from Hawthornden, v.-c. 83, in 1911, and named as above by Mr G. C. Druce.—G. C. BROWN.

*Silene inflata* Sm. Sandscale, Dalton, v.-c. 69b, July 7, 1914. The plants produced very poor capsules and no seeds. The petals were markedly cream-coloured until they were reaching maturity.—D. LUMB. "A remarkable form; I have nothing like it, excepting a plant which I gathered last July on bushy shingle near Seaton, South Devon."—E. S. MARSHALL. "Conf. *S. vesicaria* Schrad., var. *pubescens* DC., sub. var. *parvifolia* Rouy."—J. A. WHELDON.

*Silene conica* L. Roadside, dry sandy soil, Hilborough, v.-c. 28, June 11, 1914.—F. ROBINSON. Also from sandy land, Cockley Clay, 3 miles S.W. of Swaffham, W. Norfolk, v.-c. 28, June 23, 1914.—J. E. LITTLE.

*Silene anglica* L., b. *quinquevulnera* L. Hay field, Cromer, v.-c. 27, June 13, 1914.—F. ROBINSON. "Good *S. quinquevulnera* L. This seems to me much nearer to *S. gallica* L. than to *S. anglica* L., of which I cannot reckon it as a variety."—E. S. MARSHALL. "Yes, I had no previous record for the county."—G. C. DRUCE.

*Silene Otites* Wibel. Dry heath land, Barnham Common, W Suffolk, June 1, 1914.—F. ROBINSON.

*Silene nutans* L., var. *dubia* (Herbich) Williams Mon. *Silene* in *Journ. Linn. Soc.*, vol. xxxii., p. 171 (1896). Shingle, Lydd, Kent, in great quantity, July 1904. This appears to have been first described by Schur as *S. transsylvanica* in *Oester. Bot. Zeit.* viii. (1858), pp. 22 et 287. Herbich's *dubia* was published in his *Flora Bucovina*, p. 388, (1859). In the *Kew Index* both names are merged into *S. nutans*, but the publication of the latter wrongly cited as ex Rohrbach's *Monograph* of 1868, and the date as usual is suppressed. Dr Williams in his valuable *Monograph* (*l.c.*) put *dubia* as a variety of *S. nutans*, and in the same year Rouy and Foucaud (*Fl. Fr.* iii., p. 144) cites *S. dubia* and *S. transsylvanica* as synonyms of their variety *subverticillaris*, the description of which does not seem to happily fit our Kentish plant, which Mr C. E. Salmon in 1905 first clearly showed was distinct from *S. nutans*. The longer petioled and narrow, lanceolate-acute stem leaves, which are not so strongly viscid as in *S. nutans*, and the narrower and more cylindrical calyx, are marks which he rightly emphasises. If kept as a species, it should stand as *S. transsylvanica* Schur; if a sub-species, the authority is Nyman *Consp.*; if a variety, as in my *British Plant List*, then as *S. nutans* L., var. *transsylvanica*, comb. nov. A red flowered form was still earlier described as a species by Vest in *Flora* (1821), p. 50, as *S. rubens*.—G. C. DRUCE. "Yes, this

is the more graceful, less hairy, flavescent petalled plant I recorded as *S. dubia* Herb. in *Journ. Bot.* 1905, p. 127, where it is mentioned by my brother and I saw it in this station (Dungeness) in 1888.—C. E. SALMON.

*Cucubalus baccifer* L. Wood, Merton, v.-c. 28, August 6, 1914.—F. ROBINSON. "Yes, a most interesting-plant of which we know no existing station in Britain."—G. C. DRUCE.

*Cerastium nigrescens* Edmonston (= *C. arcticum* Lange). [Ref. No. 2355]. Ben Nevis, August 1903.—G. C. DRUCE.

*Cerastium vulgatum* L., var. *pentandrum* Syme. Sands of Barry, v.-c. 90, April 26, 1914.—R. & M. CORSTORPHINE. "I believe that this is var. *pentandrum* Syme (under *C. triviale* Link), though the sepals are as strongly hyaline-bordered as in *C. semidecandrum* L."—E. S. MARSHALL.

*Cerastium semidecandrum* L. Sandhills, Askham, v.-c. 69b, March 30, 1914. Is this the typical plant?—D. LUMB. "Yes, a small state."—E. S. MARSHALL. "Correctly named."—C. E. SALMON. "Yes, and a new county record for 69b."—G. C. DRUCE.

*Cerastium semidecandrum* L. St Ouen's, Jersey, April 1907. Cf. var. *congestum* Gren.—G. C. DRUCE. "This var. is not mentioned by Rouy and Foucaud. The plant seems to come under var. *glandulosum* Koch."—E. S. MARSHALL.

*Cerastium tetrandrum* Curt. Sandhills, Askham, v.-c. 69b, March 30, 1914. Ironworks, sandhills, and golf links at Askham; limestone quarry at Staunton. Many plants are to be found flowering during the first week in March. Most of the flowers seem abnormally large through their being five-parted. The lower leaves are markedly spatulate and deeply tinged with reddish purple. Mr Druce agrees with me that it is most likely *tetrandrum*, and adds the remark that in all probability much of what is named *pentandrum* is this plant.—D. LUMB. "Doubtless correct, but gathered too young and depauperate."—E. S. MARSHALL. "Right."—C. E. SALMON.

*Cerastium*—? [Ref. No. 32]. Hedgebanks, Rocklands, v.-c. 28, May 4, 1914.—F. ROBINSON. "This seems to be a robust or shade-grown form of *C. arvense* L. It agrees with the description of the var. *latifolium* Fenzl in Ledeb. *Fl. Ross.* i., p. 412 (1842), which is as follows:—'Foliis caulinis majoribus, praesertim superioribus e basi late ovata oblongis v. lanceolatis; ramorum ac fasciculorum anguste lanceolatis v. linearibus; omnibus utrinque pubescentibus, cauliculis palmaribus spithamaeis et altioribus, petalorum lobis late ovatis.'

There are specimens like this in the British Museum from one or two English localities, and also from the Continent."—A. B. JACKSON. "This must be an *arvense* form, and it has the uppermost stem-leaves broad-based, one of the characters, according to Rouy and Foucaud (*Fl. Fr.*) of var. *latifolium* Fenzl, but I have no authentic examples to compare. I have never seen this beautiful luxuriant form in Surrey."—C. E. SALMON. "Ripe fruit is not available in these examples, but *C. arvense* L., var. *latifolium* Fenzl (*C. grandiflorum* Gilib.) is like this, a tall form with broad leaves and large flowers, but I have seen no example."—J. A. WHELDON. "This is a rather notable form of *C. arvense* Linn., and appears to agree very well with the description of *C. arvense* L., var. *latifolium* Fenzl, in Rouy and Foucaud *Fl. de France* iii., p. 203. The description given of this var. is as follows:—"Feuilles caulinaires, surtout les supérieures à base large, ovales-oblongues ou sub-lancéolées celles des fascicules et des rameaux étroitement lancéolées, toutes pubescentes ou poilues sur les deux pages; tiges assez élevées; pétales à lobes ovales."—C. E. BRITTON. "An extraordinary plant, which I think comes under *C. arvense* L. It comes nearest, of the vars. given in Rouy and Foucaud, to their *c. latifolium* Fenzl, but the stems, pedicels, and calyces are densely glandular."—E. S. MARSHALL.

*Stellaria neglecta* Weihe. Hedgebank by thicket, near Great Western Station, Malvern Wells, Worcester, v.-c. 37, May 11, 1914.—C. WATERFALL. "Yes, the variation with stalked glands on the inflorescence, which I have called forma *glandulosa*."—E. S. MARSHALL. "*S. media*, var. *neglecta* Weihe, the plant with hairy pedicels and acutely tubercled seeds."—G. C. DRUCE.

*Arenaria gothica* Fr. Origin, Moughton Fell, v.-c. 64, altitude 1600 feet. A new station, where I counted fifty plants. Cultivated in my garden, Shipley, July 1914.—J. CRYER.

*Arenaria tenuifolia* L. Walls of Abbey ruins, Castleacre, v.-c. 28, May 28, 1914.—F. ROBINSON. "Yes."—E. S. MARSHALL.

*Arenaria tenuifolia* L., var. *laxa* (Jord.)? Coarse ballast, Great Northern Railway, Grove Mill, Hitchin, Herts, v.-c. 20, July 1914.—In habit these plants agree with the set from Welbury. I could find no glandular plants among them.—J. E. LITTLE. "Yes, a strong form."—E. S. MARSHALL. "My plants are almost glabrous, and do not agree with my Lakenhead *laxa*."—G. C. DRUCE.

*Arenaria tenuifolia* L., var. *laxa* (Jord.) (Det. C. E. Salmon). Hitchin, Hexton Road, near turning to Welbury, Herts, v.-c. 20, July 31, 1914.—J. E. LITTLE. "Glands very few on my two plants."—E. S. MARSHALL.

*Sagina nodosa* (L.) Fenzl, var. *glandulosa* Bess. [Ref. No. 76]. Damp part of over year hay field, Tottington, v.-c. 28, July 23, 1914.—F. ROBINSON. "*S. nodosa*, var. *viscidula* Coss. & Germ."—J. A. WHELDON. "Apparently correct; Rouy and Foucaud make it a sub-variety, perhaps rightly. Corbière seems to be the authority, under *Sagina*."—E. S. MARSHALL. Also [Ref. No. 77] from shore of Lake, Westmere, Tottington, v.-c. 28, July 24, 1914.—F. ROBINSON. "A different and much more leafy form."—E. S. MARSHALL.

*Sagina nodosa* Fenzl, var. *glandulosa* Bess. [Ref. No. 115]. Avebury Down, N. Wilts, v.-c. 7, August 28, 1914. Has this been tested by cultivation? I have distinctly glandular forms from Guernsey and Dog's Bay, Galway, and plants from Glencar, Sligo, which I thought to be quite glabrous, show one or two glandular hairs under a strong glass. It seems probable that the glandular hairs are developed only on sandy soil or in exposed situations. Similarly var. *moniliformis* Lange, to which these plants might be referred, appears to be a state of poor soil.—W. C. BARTON. "I believe so."—E. S. MARSHALL. "Yes, the Avebury Down plant and Mr Robinson's No. 76 would doubtless develop into the so-called var. *moniliformis*."—G. C. DRUCE.

*Sagina ciliata* Fr. Sandy cart ruts, Shouldham, W. Norfolk, v.-c. 28, June 24, 1914.—J. E. LITTLE. "The *Sagina* is rather puzzling, and *S. Reuteri* must always nowadays be reckoned with! However, I think you are right in calling your plant *S. ciliata*. I see the awned sepals present, which I believe are never found in *S. Reuteri*. The capsule, when over ripe, does not seem to open ultimately in the form of a cross, so *S. apetala* is ruled out."—C. E. SALMON *in lit.* "Exactly *S. ciliata* Fr., as described by Babington in the *Manual*, which has the outer sepals more acuminate and recurved, and the plant is nearly glabrous. It is according to *S. Reuteri* Lojac, but not of Boissier. This typical *S. ciliata* is very rare, if found at all, in Lancashire, and I believe the restricted plant has a south and easterly range. Dr Moss refers nearly all the plants we call *S. Reuteri* Boiss. to *S. ciliata*, and I believe favours the reduction of *S. ciliata* to the status of a variety under *S. apetala*. Whatever view is taken, I feel sure *ciliata* and *Reuteri* represent two frequently recurring extremes usually readily separable, and I think both are distinct from *S. apetala*."—J. A. WHELDON. "I believe so."—E. S. MARSHALL.

*Sagina ciliata* Fr., *forma*. [Ref. No. 52]. Cliffs near Arbroath, v.-c. 90, August 15, 1914. Outer sepals have a very short mucro, but all the sepals are obtuse and incurved at the tip like *Reuteri*; also they are much shorter than the capsule. The valves of the capsule are truncate at the tip.—R. & M. CORSTORPHINE. "A difficult plant. Very like *S. Reuteri* in habit, but I see that the outer sepals are mostly

mucronate, which is a character of *S. ciliata*. It seems to be therefore a small glandular var. of the latter."—E. S. MARSHALL. "I cannot separate this from *S. Reuteri* Boiss. The mucronate outer sepals do not recurve, and are very different in outline from the narrow acuminate ones of typical *ciliata*, as shown in Mr Little's specimens referred to above."—J. A. WHELDON. "This interesting *ciliata* form will come, I think, under var. *minor* Rouy and Fouc. (*Fl. Fr.* iii., p. 289, 1896). The whole plant is more or less glandular, but the leaves are scarcely strongly enough ciliate to agree well with Corbière's description of var. *filicaulis* Jord."—C. E. SALMON. "Nearest to *S. apetala*, var. *ciliata* Garcke (= *S. ciliata* Fries). Benekin (*Bot. Zeit.* iii., p. 721, 1845) maintained that the restricted *S. apetala* and *S. ciliata* were mere habitat states. His views were combated by Babington (*Bot. Gaz.* i., pp. 174-177, 1849), and supported by Henfrey (*Bot. Gaz.* ii., p. 182, 1850). So far as, I can judge, both *S. apetala*, var. *ciliata*, and *S. apetala*, var. *reuteri* H. & J. Groves, have glandular and eglandular forms."—C. E. MOSS.

*Sagina apetala* Ard., b. *prostrata* Gibs. St Ippolyts, Hitchin, Herts, v.-c. 20, May 29, 1913. Det. S. H. BICKHAM.—J. E. LITTLE. "Yes."—E. S. MARSHALL. "Mr Little's Herts specimens are *apetala*, but are very lax and large for *prostrata* (see *Phyt.* i., p. 178)."—G. C. DRUCE.

*Sagina apetala* Ard., var. *prostrata* Gibs. With the type, on the mud path of a newly made road near the sea, Penarth, v.-c. 41, July 7, 1913. I suppose this is correct, but it may be *S. Reuteri* Boiss., a plant which I do not know. The relative length of sepal and capsule varies (? entirely according to age) from about 3—4 to 1—2.—H. J. RIDDELSDELL. "Is not this *S. Reuteri* Boiss.? Flowers small; sepals appressed in fruit; pedicels short, or shortish."—E. S. MARSHALL. "Why not *S. ciliata* Fries, var. *ambigua* Corbière. Most of the peduncles are glabrous. The sepals are not spreading as in *apetala*."—G. C. DRUCE.

*Sagina apetala* Ard., var. *barbata* Fenzl. Sand dunes, Ainsdale, S. Lanes, v.-c. 59, July 20, 1914.—J. A. WHELDON. "Correctly named, I believe."—C. E. SALMON. "Yes, what we regard as type, I think."—E. S. MARSHALL. "Yes, I look upon this as the type with var. b. *glaberrima* Koch, = *imberbis* Fenzl, as the rarer form."—G. C. DRUCE.

*Sagina*—? Cliffs\* at Boddin, near Arbròath, v.-c. 90, June 9, 1914. This plant does not agree with any of the described forms of *S. maritima* or *S. apetala*. It grows on limestone, associated with *S. maritima* (type) and *Cochlearia groenlandica*; but it is distinguished at a glance from the former by its glaucous-grey colour.

—R. & M. CORSTORPHINE. The Rev. E. S. Marshall, to whom two batches of fresh plants were sent, writes:—"The *Sagina* strikes me as being of special interest. It clearly belongs to *S. maritima* Don. It seems nearest to var. *densa* in habit, but much less crowded; from that it also differs by the capsules exceeding the sepals. Besides this, it has some of the stems and pedicels furnished with gland-tipped hairs, which I never saw before in this species. Like var. *debilis* (Jord.), it is quite prostrate; but that is a slender spreading plant, with long internodes. Also var. *prostrata* Townsend MS. (never described, I fancy), is a plant two or three times as large and clearly different . . . The better examples of the greyish-glandular plant, in more advanced condition, still quite puzzle me. The leaves resemble *maritima* in outline; with, however, a small mucro or apiculus at the top, as a rule. The sepals are as in *maritima*; broader than in *apetala*. *S. maritima* does not appear to be ever grey or glaucous, or at all hairy or glandular. So it seems to be either a new species (at least for Britain), or a new and very marked variety of *maritima*."—E. S. MARSHALL, in *lit.* "Apparently identical with the plant sent as var. *prostrata* Towns. (Travis and Wheldon, *Report* 1913, p. 459), since identified as *S. maritima*, var. *ciliata* Nordst. We have seen an authentic specimen of Townsend's plant, which differs in being more glabrous and larger in all its parts. It has also a different habit. In comparing R. & M. Corstorphine's plant with that from South Lancashire, it must be noted that the latter grew near docks where coal is constantly loaded. They have therefore an unnaturally dark appearance."—J. A. WHELDON. "This is very interesting. Clearly, I think, a *maritima* form, which I have never seen before. There is a var. *ciliata* Nordst. in *Hartm. Skand. Fl.* 1879, p. 247, and a var. *glauca* Strobl. in *Oest. Bot. Zeit.* xxxv., 1885, p. 209, but unfortunately I have no examples of either."—C. E. SALMON. "This is undoubtedly *S. maritima* Don. I have not my specimens available at the moment, but, speaking from memory, I think the Arbroath plant is very like the *S. maritima* from Garston, distributed by Mr Wheldon and me last year, except that the former is more glandular. Our Garston plant has been referred by Dr O. Nordstedt to his var. *ciliata*, and I have no doubt that the Arbroath plant also comes under the same variety, which is probably a northern form."—W. G. TRAVIS. "An interesting plant which Mr and Mrs Corstorphine showed me *in situ* last June. The large capsules and fruiting calyx suggest *S. maritima*; but the habit, the glandular hairs, and the apiculate leaf point to *S. apetala*. Unless a hybrid, the specimens point to the desirability of uniting *S. apetala* and *S. maritima*, two plants which in any case are closely allied. It is, in my opinion, the case that the Jordanian species of *Sagina* are unduly small; on that standard one might easily make a score or two of British species of *Atriplex* and *Salicornia*. However, of the segregates or close allies of *S. apetala*, *S. maritima* is the best; but I would reduce *S. ciliata* and *S. reuteri* to varieties of *S. apetala*. I may add

that the name *S. maritima* (G. Don in *Herb. Brit. fasc. vii*, No. 155, 1806, cum descr.) is antedated by that of *S. erecta* (Müller in *Fl. Dan. fasc. 15*, p. 2, t. 845, 1782), non L., *S. erecta* L. being *Mænchia erecta*), but *S. erecta* Müller, being stillborn (since at that time *S. erecta* L. was valid), *S. maritima* Don remains the correct name for the species."—C. E. MOSS. "This is certainly not specifically distinct from *S. maritima* Don, though it does not exactly agree with any described variety. It approaches var. *densa* (Jord.), but is certainly not that form. The greyish colour ascribed to it, is probably owing to the reflection of light from the glandular-pubescent surface. The glandular hairs themselves are so minute that they do not seem to furnish a very tangible character, but taken in conjunction with the habit of the plant, the relative lengths of calyx and capsules, there are sufficient grounds to regard this plant as a well-marked var. of *S. maritima*. The apiculate leaves seem to be a varying character in this species."—C. E. BRITTON. "They are most interesting specimens. The text-books usually describe *S. maritima* as glabrous, but Rouy and Foucaud say the sepals are rarely glanduliform. These are clothed with more or less glandular hairs on the leaves, stems, and peduncles. Examination of about 200 specimens from various sources shows that specimens from Afton Bay, Isle of Wight (C. E. Palmer), have a few short glandular hairs practically confined to the peduncles; others from Headon Hill and Alum Bay, Isle of Wight (C. E. Palmer), closely approach the Scottish plant in habit, and in the hairs, which are, however, less plentiful. Specimens extremely like the Scottish plant, gathered by me at Penmon in Anglesey, were sent to the Club, but no report was made on them; a more diffuse form from Stonehaven, Kincardine, is also hairy. Specimens of var. *debilis* which I gathered at Maghera, Co. Wicklow, in 1909, have a few long ciliate hairs, especially on the leaf-sheaths. Other hairy specimens are from Mullion, Cornwall. All my other gatherings are glabrous. This Scottish plant appears to be an analogous condition to the var. *glandulosa* of *Sagina nodosa*. Mr and Mrs Corstorphine may well describe and name it."—G. C. DRUCE.

*Sagina procumbens* L., var. *apetala* Fenzl. Walton, S. Lancs, v.-c. 59, September 10, 1914. Some flowers contain a rudimentary petal, or even two, but they are mostly absent.—J. A. WHELDON. "Are such variations more than forms?"—E. S. MARSHALL.

*Claytonia perfoliata* L. Cultivated land, Walton, v.-c. 28, May 18, 1914.—F. ROBINSON.

*Montia fontana* L. = *M. lamprosperma* Chamisso. [Ref. No. 2358]. Growing on the shingle of a garden path, Glen Brittle, Skye, June 1910, showing that *lamprosperma* has both the land and water form.—G. C. DRUCE.

*Malva pusilla* Sm. Bank near Corn Mill, Ventongimps, v.-c. 1, August 11, 1914.—F. RILSTONE.

*Tilia platyphyllos* Scop. Park, Barton Seagrave, Northants, v.-c. 32, July 31 and September 5, 1914. Introduced.—G. CHESTER. "No, this is *Tilia petiolaris*, which is frequent in cultivation, but does not seem to have been found in a wild state. It is considered to be a sport of *Tilia argentea*, the White Lime of South-Eastern Europe and Asia Minor."—A. B. JACKSON.

*Tilia ulmifolia* Scop. Rockingham Woods, Northants, v.-c. 32, July 30, 1914. Rather frequent in this and other outlying woods, and probably indigenous.—G. CHESTER. "This is the small-leaved lime *Tilia cordata* Miller. The latter name has been rejected on the ground that specimen assumed to be Miller's type, in the British Museum, is *T. platyphyllos*, but as pointed out by Mr A. Henry (*Trees of Great Britain* vii., p. 1656, 1913), there is no doubt that Miller's description of *T. cordata* in the Dictionary refers to the small-leaved lime and not to *T. platyphyllos*, and his name therefore takes preference over Scopoli's *ulmifolia*."—A. B. JACKSON.

*Geranium phaeum* L. Open wood, Watton, v.-c. 28, May 17, 1914.—F. ROBINSON.

*Geranium pyrenaicum* Burm. Dry hedge bank, Croxton, v.-c. 28, June 1, 1914.—F. ROBINSON.

*Geranium rotundifolium* L. Dry hedge bank, Thetford, v.-c. 28, June 1, 1914.—F. ROBINSON. "A rare Norfolk plant."—G. C. DRUCE.

*Geranium Robertianum* L., var. *purpureum* Vill.? Shingle, Pagham, W. Sussex, v.-c. 13, June 13, 1914. "This Geranium is what we always used to call *G. purpureum*, but what it is called now I cannot say. It is a coast-shingle form, and is not, I believe, found inland." C. E. Salmon, *in lit.*, August 16, 1914.—J. E. LITTLE. "Too hairy for the plant of Foster and Villars, but still of the type undoubtedly. It is, however, what is often named *purpureum*. The carpels closely resemble those of Foster's plant. It may be the var. *rubricaulis* Horn. in Willk. et Lange's *Prod. Fl. Hisp.* iii., p. 320, but I have seen no specimens of this."—A. BENNETT. "I gathered this there in 1901. Mr Arthur Bennett wrote:—'What we so call; but Britten and Nicholson years ago denied the name.'"—E. S. MARSHALL. "Yes, the glabrous-carpelled plant under aggregate *purpureum* Vill."—G. C. DRUCE.

*Geranium Robertianum* L., var. [Ref. No. 117]. Bordeaux shingle, Guernsey, August 2, 1914.—W. C. BARTON. "Very different

from the Pagham plant, which has narrow leaflets and sub-glabrous calyces. This is very crowded and compact; leaflets broad, thin; calyces and pedicels with many long, slender, gland-tipped white hairs. I cannot name it.—E. S. MARSHALL. "This is aggregate *purpureum* Vill."—G. C. DRUCE.

*Erodium cicutarium* L'Hérit., var. *pimpinellifolium* Cav. In great quantity in a cornfield, Wigginton Heath, Oxon, July 29, 1914. Petals, 3 longer and paler; 2 shorter, darker, and spotted. Flowers vary in size. The later flowers do not show spots (? in every case). The spots consist of small very pale areas, which are, except for a narrow rim, filled with close-set very dark crimson splashes and dots, which lie in lines radiating from the very base of the petals, where the veins of the petals begin to diverge. I have never seen the spots described before, and did not know what to expect. The petals are unequal in about the proportion 2:3. The flowers are usually considerably larger than in the common plant, which abounds in such places as sand-dunes and waste ground. The var. has been found in Oxford before.—H. J. RIDDELSDELL. "Just what I understand by that."—E. S. MARSHALL. "This has the facies of Sibthorp's *pimpinellifolium*, but the petals should be spotted; an evanescent character, for I have seen spotted and unspotted flowers on the same plant. Kirscher's *Flora von Stuttgart* has a lengthy disquisition on the biology of this interesting form, which Dillenius was the first to distinguish."—G. C. DRUCE.

*Impatiens Noli-tangere* L. Woods about Stock Gill Force, Amble-side, v.-c. 69, July 27, 1914.—C. WATERFALL.

*Rhamnus catharticus* L. On carboniferous limestone, Silverdale, v.-c. 60, altitude 30 feet, May 29, 1912. These specimens were gathered from a shrub eight to nine feet high, growing by the side of the road leading to Silverdale, and almost opposite the end of the Carnforth road. The bush has since been cut down to widen the road. The yellow-green colour and mealy character of the foliage were very striking when first gathered.—J. CRYER.

*Acer campestre* L., var. *leiocarpon* Wallr. Ashmansworth, N. Hants, v.-c. 12, September 18, 1914. All these specimens have been seen by Mr A. B. Jackson who assents to the naming.—W. C. BARTON.

*Medicago Falcata* L. Among gorse on dry heath, Barnham Common, v.-c. 28, August 17, 1914.—F. ROBINSON. "Yes."—E. S. MARSHALL. "Nice specimens from, I presume, a native station which is not mentioned in Nicholson's new *Flora of Norfolk*."—C. E. SALMON.

*Medicago sylvestris* Fr. Roadside, gravelly soil, Bawburgh, v.-c. 27, July 3, 1914.—F. ROBINSON.

*Medicago sylvestris* Fr., var. *procumbens* Fr. = *M. cyclocarpa* Hy. [Ref. No. 1352]. Between Mickleham and West Humble, Surrey, August 23, 1914. Growing on the border of a field cropped with lucerne. I have not been able to see any material quite like my plant in the collections at the Natural History Museum, South Kensington. My plant agrees with Fries' description of his variety, and also with the plant called by Rouy *M. sylvestris* Fries, var. *cyclocarpa* Hy. As will be seen, the distinguishing features are the long diffuse stems, narrow leaflets, the dusky yellowish-purple corollas, and the ring-like legumes.—C. E. BRITTON. "I think this a *M. falcata* × *sativa*, nearer to *sativa* in flowers. Young pods with many appressed silky hairs."—E. S. MARSHALL.

*Medicago minima* Desr. Dry hillside, chalk soil, Hillborough, v.-c. 28, June 11, 1914.—F. ROBINSON.

*Medicago lupulina* L., var. *Willdenowiana* Koch? Roadside, Guildford, Surrey, v.-c. 17, September 1914.—J. COMBER. "No doubt the hairs on the pods are glandular, but one can hardly call them 'yellowish.'"—A. BENNETT. "Yes, fruit glandular. The habit is that of cultivated *M. lupulina*, in my specimens."—E. S. MARSHALL. "Pods with glandular hairs, i.e., var. *Willdenowiana* Koch."—C. E. SALMON.

*Medicago lupulina* L., var. *eriocarpa* Rouy. Formby, S. Lanes, v.-c. 59, July 15, 1914.—W. G. TRAVIS. "Agrees with the description, but Rouy makes it a sub-variety."—E. S. MARSHALL. "Strictly speaking, it is sub-var. *eriocarpa* Rouy. 'Légume pubescent ou velu; plante ordinairement fortement pubescente-soyeuse,' and this plant only approaches these characters."—G. C. DRUCE.

*Trifolium scabrum* L. Dry hillside, chalk soil, Hillborough, v.-c. 28, June 11, 1914.—F. ROBINSON.

*Trifolium subterraneum* L. Heath land, near water, by Punch Bowl, Croxton, v.-c. 28, May 3, 1914.—F. ROBINSON. Also from Lancut, v.-c. 34, June 12, 1913. A rare plant in W. Gloster; this is a new locality for it.—H. J. RIDDELSDELL. Also from Colney Heath, Herts, v.-c. 20, May 19, 1913.—J. E. LITTLE.

*Trifolium minus* Relh., var. *microphyllum* Ser. Sandhills, Hightown, S. Lanes, v.-c. 69, July 7, 1914.—W. G. TRAVIS.

*Trifolium dubium* Sibth., var. *pygmaeum* Soy.-Will. Foot of cliffs, Arbroath, August 15, 1914.—R. & M. CORSTORPINE. "According to Rouy, this variety was described under the name *T. filiforme*, and is the same as *T. minus* Relhan, var. *microphyllum* Seringe, in

DC. *Prodrömus*. The Arbroath cliff plant I can only look upon as starved type."—E. S. MARSHALL. "Yes, the correct name for the species is *T. dubium* Sibth. Fl. Ox. 1794."—G. C. DRUCE.

*Securigera Securidaca* (L.). Garden weed, Dalton, v.-c. 69b, August 9, 1914. If attempts be made to grow this from the seeds supplied, the testa should be pierced before sowing, otherwise most of them will remain like a piece of flint, no matter how wet the soil may be. It is a most interesting plant to watch. The earliest umbels, in this quarter at anyrate, are microscopic in almost every particular. Mr Druce kindly named the plant for me. The faded unpressed pieces are included to show the length of the mature umbel stalks.—D. LUMB. "Yes, the *Bonaveria Securidaca* (L.) Desv. The genus *Bonaveria* Scopoli dates from 1777; that of *Securigera* DC. only from 1805."—G. C. DRUCE. "Yes."—A. THELLUNG.

*Lotus siliquosus* L. (*Tetragonolobus siliquosus* Roth). Chalk Downs, near Streatley, Berks, May 1914. First found 1911, well established.—Coll. V. C. MURRAY; comm. G. C. DRUCE.

*Lotus uliginosus* Schk., var. *glabriusculus* Bab. Wet lane, Petit Bot, Guernsey, August 1, 1914. Very similar to a plant distributed last year through the Watson Exchange Club by Mr Standen, and so named by Mr Salmon. Is it var. *glaber* of Bréb., and the same as var. *a. sub-glaber* of Syme E.B.—"sub-glabrous with the leaflets ciliated at the margins?" These specimens grew in the damp hedgerow of a "water lane" and were very different in general appearance from the hairy form growing near by on dryer ground.—W. C. BARTON. "Doubtless right. But I think that the amount of hair varies according as the locality is wet or dry."—E. S. MARSHALL. "May pass, I think, but not so extreme as specimens sent me by Mr R. S. Standen from Lindfield, Sussex, in 1911. Rouy (*Fl. Fr.*) segregates *L. uliginosus* into *L. glabriusculus* Bab. and *β. villosus* Lamotte. Our commoner plant is certainly the latter. Rouy remarks that the flowers of *glabriuscula* do not become green (or only slightly so) when dry, whereas those of *villosus* usually show this change. This is borne out in my herbarium specimens and in Mr Barton's example before me."—C. E. SALMON. "Yes, but Bab. in ed. 2 of his *Manual*, put it as a variety of *L. major*, hence if his name for it is used 'Bab.' should be in brackets. Brébisson in his *Fl. Norm.* 87 (1858), under *L. uliginosus* had a var. *glaber* (as in my *List*)."—G. C. DRUCE.

*Lotus tenuis* Waldst and Kit. Rough pasture, clay washing on chalk, under "Eagle's Nest," Offley Hill, Hitchin, Herts, v.-c. 20, August 10, 1914. *L. tenuis* in N. Herts occurs (1) on heavy boulder clay; (2) on the chalk scarp, in poor soil with a washing of marl or of clay from the caps on the hills. It generally occurs with *L. cornicu-*

*latus*, and these appear to be intermediate forms. At Pagham, W. Sussex, v.-c. 13, it occurs on the sea bank in apparently drier situations, though perhaps moisture soaks up from below.—J. E. LITTLE. "Very characteristic."—E. S. MARSHALL.

*Astragalus glycyphyllos* L. Hedgebank, Holme Hall, v.-c. 28, August 26, 1914.—F. ROBINSON.

*Astragalus danicus* Retz. Newmarket Heath, Cambridge, v.-c. 29, June 8, 1913.—W. C. BARTON. Also sent from dry heath, near Icklingham, W. Suffolk, June 4, 1914.—F. ROBINSON.

*Oxytropis sericea* (Lam.) Simonk. Bettyhill, Sutherland, July 1907. Growing very plentifully in blown sand on the coast of West Sutherland at Bettyhill. It was in splendid flower, and was a conspicuous object in the flora of the district. The name *uralensis* must, however, give way to that of *O. sericea* Simonkai, the trivial of which is earlier than *uralensis*.—G. C. DRUCE.

*Vicia lutea* L. Beacon Hill, St Osyth, N. Essex, v.-c. 19, July 5, 1914.—G. C. BROWN. Also sent from field near Lower Sydenham, W. Kent, June 1912. Introduced.—Coll. H. B. FOXLEE; comm. J. GROVES.

*Vicia Lathyroides* L. Maulden, Beds, v.-c. 30, April 25, 1914. Many of the roots have come up uninjured from the light sand and show nodules, which I suppose to be *Rhizobia* containing bacteria and assisting the plant to obtain a supply of nitrogen.—J. E. LITTLE.

*Lathyrus palustris* L. Marsh land, edge of ditch, Woodbastwick, v.-c. 27, July 9, 1914.—F. ROBINSON.

*Rubus plicatus* Wh. & N., *forma*. Bognor Common, Fittleworth, W. Sussex, v.-c. 13, July 1914.—L. CUMMING. "A very singular and constant little form. Though strongly recalling *R. fissus* in prickles and stem pieces, it goes better under *R. plicatus*, even its prickles and stem leaves suiting *plicatus* better. Small and slender as the prickles are, they are really too few in number, too strong, and often too stout based for *fissus*; while the basal leaflets of the stem leaves are not always strictly sessile and so hardly differing at all from those of *plicatus* in July."—W. M. ROGERS.

*Rubus affinis* Wh. & N., var. *Briggsianus* Rogers. Near St David's, Pembroke, August 20, 1903. See *Report* 1903, p. 14.—Coll. AUGUSTIN LEY; comm. G. C. DRUCE.

*Rubus imbricatus* Hort. Glen Frome, near Stapleton, Gloucestershire, August 20, 1913.—J. W. WHITE.

*Rubus sciaphilus* Lange. Grwyne Valley, Brecon, August 13, 1903. See *Report* 1903, p. 15.—Coll. AUGUSTIN LEY; comm. G. C. DRUCE.

*Rubus thyrsoides* Wimm. Wyre Forest, Shropshire, September 15, 1903. See *Report* 1903, p. 15.—Coll. AUGUSTIN LEY; comm. G. C. DRUCE. "Not *R. thyrsoides* Wimm., I feel sure; but a good form of the variable *R. argentatus* P. J. Muell."—W. M. ROGERS.

*Rubus robustus* P. J. Muell, *forma*. Badby Wood, Northants, v.-c. 32, July 1913 and 1914.—L. CUMMING. "I see no reason for separating this from *R. robustus*, though the long lax pyramidal panicles are somewhat abnormal. It seems best to let the name cover a series of forms, as it does now with us—off type towards my *subinermis*."—W. M. ROGERS.

*Rubus silvaticus* Wh. & N. Badby Wood, Northants, v.-c. 32, July 1913.—L. CUMMING. "As far as I know, a record for Northants."—W. M. ROGERS.

*Rubus lentiginosus* Lees. Fittleworth, W. Sussex, v.-c. 13, July 1914.—L. CUMMING.

*Rubus macrophyllus* Wh. & N., *forma*. Badby Wood, Northants, v.-c. 32, July 1914.—L. CUMMING. "A hairy form towards *Schlechtendalii*."—W. M. ROGERS.

*Rubus adenanthus* Boul. et Gill. Below Peckforton Castle, Cheshire, August 5, 1903. See *Report* 1903, p. 17. Ex herb. A. H. WOLLEY-DOD; comm. G. C. DRUCE. "Apparently."—W. M. ROGERS.

*Rubus anglosaxonicus* Gel., var. *vestitiiformis* Rogers. Woods at Ross, Herefordshire, September 5, 1903.—Coll. AUGUSTIN LEY; comm. G. C. DRUCE.

*Rubus Drejeri* G. Jensen. Durdham Down, Bristol, W. Glos., August 1, 1914.—J. W. WHITE. "I agree."—W. M. ROGERS.

*Rubus Babingtonii* Bell Salter. Common round Fittleworth, W. Sussex, v.-c. 13, July 14, 1914.—L. CUMMING.

*Rubus fuscus* Wh. & N. Bognor Common, and elsewhere near Fittleworth, W. Sussex, July 1914.—L. CUMMING.

*Rubus fuscus* Wh. & N., var. *macrostachys* P. J. Muell. Coldborough, Herefordshire, September 8, 1903.—Coll. AUGUSTIN LEY; comm. G. C. DRUCE.

*Rubus scaber* Wh. & N.? Badby Wood, Northants, v.-c. 32, July 1914.—L. CUMMING. "Apparently *R. scaber*, but, if so, with glandular development on stem abnormally weak. *R. scaber* would be a record for Northants."—W. M. ROGERS.

*Rubus glareosus* Rogers, nov. sp. Bognor common and other stations in W. Sussex, v.-c. 13, July 1914.—L. CUMMING.

*Rubus Marshalli* Focke & Rogers. Common near Fittleworth, W. Sussex, v.-c. 13, July 1914.—L. CUMMING.

*Rubus Kaltenbachii* Metsch. Newent wood, W. Gloucester, September 10, 1903.—Coll. AUGUSTIN LEY; comm. G. C. DRUCE.

*Rubus hirtus* Waldst. & Kit., var. *flaccidifolius* *a* and *b*. Badby Wood, Northants, v.-c. 32, July 1914.—L. CUMMING. "There seems no reason for keeping forms *a* and *b* apart; though *b* sheets seem on the whole further away from *flaccidifolius* towards *rotundifolius*. But I have to own (with these packets before me) it may be open to question whether these two varieties can profitably be kept apart. Both varieties are records for Northants."—W. M. ROGERS.

*Rubus nutkanus* Mocino. Woods near Colliston, v.-c. 90, May 28, 1914. See *B.E.C. Report* 1913, p. 316.—R. & M. CORSTORPHINE.

*Rubus*—? Naturalised in several stretches of wood near Leysmill, Forfar, v.-c. 90, July 3, 1914.—R. & M. CORSTORPHINE. "This reminds me of *R. spectabilis* Pursh, naturalised near Sandling Park, S. Kent, and locally called (*vide* Dr Cosmo Melvill) 'the woodman's rose.' I have gathered it, but cannot find my specimen."—E. S. MARSHALL.

*Potentilla thuringiaca* Bernh., var. *Nestleriana* Schinz and Keller. Railway bank near Friockheim, Forfar, v.-c. 90, July 3, 1914. See *B.E.C. Report* 1910, p. 500.—R. & M. CORSTORPHINE.

*Potentilla verna* L. Limestone crags near Wynd's Point, Hereford, v.-c. 36, May 11, 1914.—Coll. R. F. TOWNDROW; comm. C. WATERFALL. Also sent from Carboniferous Limestone, Silverdale, v.-c. 60, altitude 30 feet, May 30, 1914. Occurs in several stations in and around Silverdale. This year it was seen to great advantage in a rough and rocky pasture leading towards Silverdale Moss.—J. CRYER.

*Potentilla procumbens* × *erecta* = *P. suberecta* Zimm. Peat moor near Ashcott Station, N. Somerset, August 6, 1914.—J. W. WHITE. "Yes, a slender heath-form of the hybrid, which I have seen near Shapwick station, not far away."—E. S. MARSHALL.

*Alchemilla acutidens* Bus., var. *alpestriformis* C. E. S. Origin near Lochan-na-Chat, Ben Lawers, 1913. Hort. Reigate, Surrey, Aug. 1914. See *Journ. Bot.* 1914, p. 287. It was noticeable how much smaller these plants kept compared with *A. alpestris* grown alongside in the garden.—C. E. SALMON.

*Rosa sempervirens* L., b. *Melvini* (Towndrow). Edge of pond, Leigh Linton, Worcester, v.-c. 37, May 7, 1914.—Coll. R. F. TOWNDROW; comm. C. WATERFALL. "So very much off *R. sempervirens*, as I have seen it in N. France, that I incline to think it specifically distinct. Not known elsewhere, I believe."—E. S. MARSHALL.

*Rosa canina* L., agg. [Ref. No. 4951.] Hanwell, Oxon, Aug. 1910.—G. C. DRUCE. Comes between (l) *andegavensis* Bast. and (p) *verticillacantha* Mérat; that is, some leaflets have most or all of the teeth simple, whilst others have them more or less compound. According to Keller and Christ, *R. hirtella* Rip. is a somewhat similar form, but with oval fruit and sepals glandular on back."—W. BARCLAY.

*Rosa dumetorum* Thuill., var. *platyphylla* Rau. [Ref. No. 3.] Hedge, Grey Abbey, Co. Down, Sept. 3, 1914. Leaves irregular in size; some very large, very dark green above. Approaching biserrate towards top of leaf. Fruit single; some very large and turbinate.—C. H. WADDELL. "This is a form of *R. dumetorum* Thuill. The (g) *platyphylla* Rau., so far as I can make out, differs from (d) *urbica* chiefly by having leaflets broader in proportion to their length. The present specimen does not show this difference. It had best be set down as a glaucous form of (d) *urbica*, with globose instead of ovoid fruit."—W. BARCLAY. "I do not think this can be *R. platyphylla* Rau., which, according to descriptions, is a plant with much broader leaflets—orbicular or very broadly oval—and fruit tending to a more ovoid form. I should call this *R. sphaerocarpa* Pug."—C. E. BRITTON.

*Rosa glauca* Vill. [Ref. No. 1]. Grey Abbey, Co. Down, September 3, 1914.—C. H. WADDELL. "Not materially different from No. 2. Serration rather more compound. Both are glaucous forms, and might therefore, according to Baker, come under var. *glaucophylla* Winch."—W. BARCLAY. "Fruit subglobose; sepals deciduous, patent or loosely reflexed; leaflets slightly compound-serrate. It can hardly be any form of *R. glauca*. Is it not a '*canina*' (aggregate), of the *S. sphaerica* Gren., or *R. globularis* Franchet affinity?"—E. S. MARSHALL.

*Rosa glauca* Vill. [Ref. No. 2]. Grey Abbey, Co. Down, September 3, 1914.—C. H. WADDELL. "This is not a form of *R. glauca* but *R. canina* L. (g) *dumalis* Bechst."—W. BARCLAY.

*Rosa Eglantheria* L. Peppard, Oxon, July 1906.—G. C. DRUCE.  
 “More advanced fruit necessary to say whether *R. comosa* Rip.”—  
 A. H. WOLLEY-DOD.

*Rosa suberecta* Woods, var. Westridge Wood and Nibley Knoll,  
 W. Gloucester, July 10 and September 29, 1904.—J. W. WHITE.  
 “This is doubtless a variation of the *omissa* group of *R. tomentosa*  
 Sm. It is certainly not a variety of *R. suberecta* Woods. Nor do I  
 see how it can come under *R. suberecta* Ley, as it differs in all the  
 characters relying on which that so-called species has been segregated.  
 It has not prickles straight or nearly so, but stout and decidedly falcate.  
 Its calyx tube is not densely aciculate, the fruit is not globose. The  
 petioles have not ‘numerous unequal falcate acicles and pricklets,’ but  
 are mostly quite unarmed. No red colour is perceptible in the younger  
 parts, and not more than is common to many species in the older  
 branch. It might as easily be made a variety of *R. Sherardi* Ley and  
 more easily a var. of *R. Andrezovii* Ley.”—W. BARCLAY. “Flowering  
 examples only, showing very little of the armature. I consider this  
 to be *R. tomentosa* Sm., and cannot see any likeness to Ley’s own  
 examples of his species—*R. suberecta* Ley = *R. villosa* L., var. *sub-  
 erecta* Woods.”—C. E. BRITTON.

*Rosa tomentosa* Sm. [Ref. No. 1368.] Marden Park, Surrey,  
 Sept. 6, 1914.—C. E. BRITTON. “Belongs, I think, to the  
*scabriuscula* group of *R. tomentosa* Sm., differing from var.  
*scabriuscula* Sm. in its more hairy and somewhat glaucous leaflets,  
 its prickles less slender and more hispid styles. The serration of  
 the leaves is also less deep and less compound.”—W. BARCLAY.

*Rosa pimpinellifolia* L. × *tomentosa* Sm. [Ref. No. 69 (1).]  
 Elliot Den, near Arbroath, v.-c. 90, July 27 and September 13, 1914.  
 Growing with both parents. The *tomentosa* forms near it belong to  
 the groups *subglobosa* Sm. and *omissa* Déségl. Serratures nearly  
 simple, fruit broadly ovate or almost globular, with a short neck.—  
 R. & M. CORSTORPHINE. “No. 69 (1) has a look of a *mollis* ×  
*spinosisima*, to my eye.”—E. S. MARSHALL.

*Rosa involuta* Sm., var. [Ref. No. 66 (3).] Cuthlie Den, near  
 Arbroath, v.-c. 90, July 27 and September 13, 1914. Bush  
 —tall, straggling; petals—large, pink; fruit—long, urceolate,  
 with longish neck; leaflets—almost uniserrate, slightly hairy above,  
 more so beneath, with a few glands on the midrib.—R. & M.  
 CORSTORPHINE. “An interesting hybrid. The second parent can be  
 better guessed at on the spot.”—E. S. MARSHALL. “These two forms  
 are practically the same, except that in the second the fruit is  
 elliptic-oblong, or in some cases turbinate, whereas in the former it is,  
 as is more usually the case, globose. Both have the serration

simple, except that here and there a toothlet may be detected. Both have the leaves sparingly hairy and quite eglandular, except that on the midrib an occasional gland may be detected. The main sepals in both have some slender pinnae, and in both the pedicels and fruits are pretty thickly clothed with glands and acicles. Both have glaucous (blue-green) leaves, and their prickles are similar in form. In the second the fruit as I saw it on 29th July, before it had begun to shrivel, was remarkable for its lengthened shape, ovoid prolonged or in some cases obovoid, quite different from any other form which I have seen. The mature fruit is not so striking. At some distance, but in the same locality, was another clump, somewhat similar to the first, but with more double teeth and with leaflets broadly elliptical, rounded at both ends and occasionally almost orbicular. The leaves were strongly tinged with red even at that early part of the season. Still a fourth bush or clump differed in its leaves with composite glandular serration and with numerous subfoliar glands. All four, however, appear to me to be variations of *R. pinupinellifolia* × *tomentosa* Sm. Mr and Mrs Corstorphine are to be congratulated on the discovery of these interesting forms."—W. BARCLAY.

*Rosa involuta* Sm., *forma* = *R. spinosissima* L. × *tomentosa* Sm. August 18, 1914. This is the rose described by Major Wolley-Dod in *List of British Roses*, p. 9, under the name of *R. spinosissima* (agg.) × *dumetorum* (agg.) or (*coriifolia*? agg.) f. *Margerisoni* f. nov. The specimens enclosed are from a plant in my garden, sent me about 7 years ago by Mr Margerison, who discovered it in Knipe Wood, Kettlewell, N. Yorks. I formerly sent a moderate supply to the Watson Club, and what I now send should enable all who take an interest in roses to obtain a specimen. To the note published in the 28th Annual Report of the Watson Club I have nothing to add except that I am more convinced than ever that this rose belongs, not to the *hibernica*, but to the *involuta* group.—W. BARCLAY. "I presume that this is identical with the rose described by Wolley-Dod under the cumbersome form of *R. spinosissima* (agg.) × *dumetorum* (agg.) (or *coriifolia*? agg.) f. *Margerisoni*, in his *List of British Roses*. How much more satisfactory would it have been had Wolley-Dod simply described this as × *Rosa Margerisoni*! These excellent specimens afford a better opportunity of studying this rose than the late-gathered specimens contributed by Mr Druce in 1911. That this is a *spinosissima* hybrid is, I believe, uncontroverted; the only point on which opinion is divided is as to the second parent. As to *R. tomentosa* or *R. omissa* entering into the composition of this rose, I am quite unable to believe. The presence of a *tomentosa* form would be revealed by a glandular development, whereas this rose is practically eglandular. The very narrow stipules are against the view of *R. coriifolia* being the second parent, and I think Wolley-Dod's first alternative *R. spinosissima* × *dumetorum* is the correct solution. The examples have a certain

resemblance to *R. hibernica*, but clearly cannot be ranged under that name, and, scientifically, it is impossible to call this *R. involuta* Sm., as Mr Barclay does, the points of divergence being so great."—C. E. BRITTON. "Nearer to the *spinosissima* parent than these hybrids usually are. At first glance it suggested to me *R. mollis*, var. *coerulea*, as the other parent."—E. S. MARSHALL.

*Rosa involuta*, var. *Wilsoni* Borrer. Growing on a very restricted area on a bank at the edge of the Menai Strait, near Bangor, N. Carnarvon, v.-c. 49, September 7, 1888.—C. BAILEY.

*Pyrus torminalis* Ehrh. "Long Cross," near St Newlyn East, W. Cornwall, June 1, 1914, and August 7, 1912 and 1914. Fairly good specimens, I think.—C. C. VIGURS.

*Crataegus monogyna* Jacq., forma *subtrilobata* Druce. [Ref. No. 154.] Virley, N. Essex, v.-c. 19, May 31, 1914. Teste G. C. Druce; who says *in lit.* "A very interesting form—forma *subtrilobata*."—G. C. BROWN.

*Crataegus oxyacanthoides* Thuill., var. *macrocarpa* Heger. [Ref. No. 70.] Virley, N. Essex, v.-c. 19, May 31, 1914; fruit, September 6, 1914. Teste G. C. Druce. Mr Druce says *in lit.* "The leaves are not quite typical *oxyacanthoides*. The size of the fruit brings it under var. *macrocarpa* Heger, which I have seen in Essex, both North and South."—G. C. BROWN.

*Crataegus oxyacanthoides* Thuill. (*oxyacantha* L.). Hedgerow in lane near Melmerby, Cumberland, v.-c. 70, August 3, 1914.—C. WATERFALL. "Under this, but not quite typical. Several forms have only one style and the fruit is small, but the leaf-veining is correct. I believe it is a new county record for v.-c. 70."—G. C. DRUCE.

*Crataegus oxyacantha* L., var. Single tree at Wigginton, Oxon, October 19, 1914. I doubt this identification; for though there are two styles, and the fruit is apparently smooth, yet the veins of the leaves are frequently not incurved. The remarkable feature of the tree is the flattened fruit, which is of about the same diameter as the usual *Crataegus* fruit, but very little more than half the length. The haws packed separately shew the feature. The tree was noticed too late for good specimens to be procurable. It is hoped that next year better and completer specimens may be distributed, and some name may be found for the variety.—H. J. RIDDELSDELL. "Material shabby; style solitary; so it appears to be a striking leaf-variety of *C. monogyna* Jacq."—E. S. MARSHALL. "An interesting looking plant, but the specimen is without flowers, and the fruit is imperfect.

I should like to see more and better specimens, but I expect it is forma *triloba*, unless it be a hybrid with *oxyacanthoides*."—G. C. DRUCE.

*Crataegus punctata* Jacq. Planted ground, Dalton-in-Furness, v.-c. 69b, June 15, 1914.—D. LUMB. "In the absence of fruit it is not possible to name this definitely. In any case, being a cultivated tree, it is of no interest to the members."—A. B. JACKSON. "Yes, similar to the plants so named by Dr Thellung."—G. C. DRUCE.

*Saxifraga aizoides* L. Mountain streamlet above Easedale Tarn, Westmoreland, v.-c. 69, July 19, 1914.—C. WATERFALL.

*Saxifraga hypnoides* L., agg. [Ref. No. 2902.] Ben Bulbin, Sligo, June 1909.—G. C. DRUCE. "Yes, and under the type, I should say."—E. S. MARSHALL.

*Saxifraga stellaris* L. Swamps on descent of Red Screes towards Scandale Beck, near Ambleside, Westmoreland, v.-c. 69, July 29, 1914.—C. WATERFALL.

*Saxifraga hirsuta* L., var. *acutidens* E. S. Marshall. [Ref. No. 3647.] Roots from cliffs above Lough Doon, Connor Hill, S. Kerry, June 1911; flower garden, West Monkton, May 24, 1914. The Linnean type has crenate leaf-toothing; in var. *acutidens* it is serrate. This is the usual Irish form, and varying considerably in the shape of the lamina, and in the amount of hairiness. The leaf-bases are truncate or somewhat cuneate; I believe that a cordate base indicates crossing with *S. Geum*.—E. S. MARSHALL.

*Chrysosplenium alternifolium* L. Wet places, Via Gelia, Derbyshire, v.-c. 57, April 12, 1914. Intermixed with *oppositifolium*.—G. C. CHESTER.

*Tillaea muscosa* L. Sandy cart ruts, Shouldham, W. Norfolk, v.-c. 28, June 24, 1913.—J. E. LITTLE.

*Sedum Forsterianum* Sm., type (Watson's *a. virescens*). [Ref. No. 4034.] Root from Culbone Woods, S. Somerset, v.-c. 5. After several years' cultivation its green foliage is maintained; and it is less robust than the var. *glaucescens*, from the coast near Minehead; garden, West Monkton, June 24, 1914.—E. S. MARSHALL.

*Callitriche obtusangula* Le Gall. Pools on Upper Chase Road, Malvern, Worcester, v.-c. 37, May 13, 1914.—C. WATERFALL. "I am unable to pass an opinion on such specimens."—A. BENNETT. "Neither habit nor fruit suggest this, to me; the latter is flat, not obtusely angled. I think it a form of *C. stagnalis* Scop., but more

careful drying would give one a better chance to judge."—E. S. MARSHALL. "Probably *C. stagnalis*."—G. C. DRUCE.

*Epilobium roseum* Schreb. [Ref. No. 658.] By a small spring in a clayey bank, Halstead, N. Essex, v.-c. 19, August 23, 1914.—G. C. BROWN. "No; certainly a state of *E. obscurum*."—E. S. MARSHALL.

*Epilobium alsinefolium* Vill. Mountain streamlet, the Red Screes, over Kirkstone Pass, Westmoreland, v.-c. 69, July 29, 1914.—C. WATERFALL. "Yes."—E. S. MARSHALL.

*Circaea alpina* L., b. *intermedia* Ehrh. Roadside, on way to Jenkin's Crag, Ambleside, Westmoreland, v.-c. 69, July 22, 1914.—C. WATERFALL. "I cannot separate this from the type. *C. intermedia* Ehrh. is a much larger plant."—E. S. MARSHALL.

*Astrantia major* L., var. *involuta* Koch. Tayside, Mid Perth, July 1905. Quite naturalised on the banks of the Tay below Perth, where very luxuriant plants are to be seen. The variety is described in the *Synopsis Fl. Germ.*, p. 280, 1837, as "involucris foliola umbella sesquialongiora, ad apicem rarius uno-alterove denticulo, nec vero in omnibus foliolis, neque in singulis regulariter, instructa."—G. C. DRUCE.

*Cicuta virosa* L. Banks of river, Salhouse, v.-c. 27, July 9, 1914.—F. ROBINSON.

*Pimpinella Saxifraga* L., var. *dissecta* Druce. Avebury Down, N. Wilts, v.-c. 7, August 5, 1913.—W. C. BARTON. "Towards this, but character not very good, in my one small plant."—E. S. MARSHALL.

*Anthriscus vulgaris* Bernh. Seedlings. Sandy bank, near Kettering, v.-c. 32, March 22, 1914.—G. CHESTER.

*Selinum Carvifolia* L. Chippenham Fen, Cambridge, August 1903. Of somewhat uncertain occurrence. That year it was in great profusion, but in other years but few plants were seen. Unless intentionally sown there, it has the appearance of being native.—G. C. DRUCE.

*Galium Mollugo* L., var. *Bakeri* Syme. Cliff tops, Milford-on-Sea, S. Hants, August 1914.—J. COMBER. "To me a reduced form of *Mollugo*. The leaves are unlike the var. *Bakeri*."—A. BENNETT. "Under that, I think; but some of the leaves broaden out unusually upwards."—E. S. MARSHALL. "Surely not *Bakeri* which should have more or less linear-lanceolate leaves, etc. Is not this a reduced state of ordinary *Mollugo* (*elatum*), frequently found on sea cliffs?"—C. E. SALMON.

*Valerianella dentata* Poll. Clears, Reigate, Surrey, June 30, 1914. This, at first sight, was taken for the variety *micota*, but on closer examination the fruits appeared diseased. Mr Ramsbottom kindly examined the specimens and reported that the whole plant was affected by *Erysiphe Polygoni* DC., a "mildew" that attacks a very large number of species of plants.—C. E. SALMON.

*Valerianella rimosa* Bast. Wheat field, Cromer, v.-c. 27, June 21, 1914.—F. ROBINSON. "I should name this *V. dentata* Poll., not *V. rimosa*."—C. E. SALMON. "*V. dentata*."—G. C. DRUCE.

*Filago minima* Fr. Sand dunes, Ainsdale, S. Lancs, v.-c. 59, July 25, 1914.—J. A. WHELDON. Also from sandy roadside, Freshfield, S. Lancs, v.-c. 59, July 1914.—W. G. TRAVIS. "We have been wrong in giving Fries as the authority for this name. Persoon *Syn.* ii., p. 422, 1807, is the earlier publication, Fries' *minima* not appearing till 1822. Mr Travis's specimens approach the var. *supina* Rouy and Camus *Fl. Fr.* viii., p. 176. Mr Wheldon's are nearer the var. *brevifolia* of those authors."—G. C. DRUCE.

*Gnaphalium luteo-album* L. Sandy land formerly cultivated, Thompson, v.-c. 28, July 23, 1914.—F. ROBINSON. "Nice specimens. The label describes them as occurring on 'land formerly cultivated.' Nicholson (*Fl. Norf.* 1914, p. 97) calls this species 'doubtfully native' in the county, but I think that the coast stations (where the plant occurs on apparently virgin soil) need not be regarded with much suspicion."—C. E. SALMON.

*Gnaphalium sylvaticum* L. Rough pasture between Shilling Green and Little Hill End, Herts, v.-c. 20, September 9, 1913.—J. E. LITTLE.

*Chrysanthemum*—? [Ref. No. 628]. On cotton-seed refuse, Hythe Quay, Colchester, v.-c. 19, June 1914. Near *segetum*, but flower somewhat deeper yellow and leaves bright green. Plants all small.—G. C. BROWN. "*Chrysanthemum coronarium* L."—A. THELLUNG.

*Artemisia campestris* L. Among gorse on dry heath, Barnham Common, v.-c. 28, August 17, 1914.—F. ROBINSON. "Beautifully prepared specimens. On the Continent this is a variable species, 20 varieties being described in the *Flore de France*."—G. C. DRUCE.

*Senecio viscosus* L. Railway track, Sandsend, Yorks, v.-c. 62, August 13, 1913.—J. E. LITTLE. "Yes. I have found it near Ware."—G. C. DRUCE.

*Senecio vulgaris* L., *forma*. [Ref. No. 121]. Stiff clay on dyke, Brean Down, N. Somerset, v.-c. 6, April 26, 1914.—W. C. BARTON.

*Senecio lantus* Forst. Banks of Tweed, Selkirk, v.-c. 79, September 1914. Native of Australia. Growing in this neighbourhood it does not increase by its own fruits, but by rooting all along its old wood. On a plant of two years' growth I counted 70 old stems, and the branches shooting from them each bearing on an average 190 blossoms.—I. M. HAYWARD.

*Cirsium lanceolatum* Scop.  $\times$  *acaule* Weber. Undercliff, Milford-on-Sea, S. Hants. v.-c. 11, August 1914. I send a few more sheets of this plant to supplement those sent last year. A further examination *in situ* convinces me that whatever it may prove to be, it is certainly not the typical caulescent state of *C. acaule*. The phyllaries have patent or sub-patent spinous tips, whereas those of *C. acaule* are mucronate only. The leaf segments are longer, narrower in proportion to their length, and more parallel sided; and the petioles, especially of the lower leaves, more spinous-ciliate. The whole facies of the plant is different—greyer in colour, and considerably rougher and coarser looking. It also flowers at least a week or ten days earlier.—J. COMBER. "See *Report* 1913, p. 476. I can add nothing to my remarks, except to say that one is better qualified to judge after seeing the plant growing with its supposed parents. Dr Thellung says: scarcely  $\times$  *lanceolatum*."—G. C. DRUCE.

*Cnicus oleraceus* L. Marshy meadow at Limehaugh, E. Perth, August 11, 1914. This thistle appeared first in 1911 in a marshy meadow close by the River Tay, which in very high floods is liable to be inundated. It has appeared every summer since to the number of 60 or 70 flowering heads. As the meadow is cut about the end of August, it is doubtful if the plant produces ripe seed so as to give it the chance of becoming thoroughly naturalised. Although not a native of Britain it is common in Central Europe and in Scandinavia.—W. BARCLAY. "Being found in South Scandinavia, Denmark, Holland, Belgium, France, and Germany, it should occur as a native in Britain."—E. S. MARSHALL. "Yes, *Cirsium oleraceum* Scop. Also naturalised near Selkirk (Hayward)."—G. C. DRUCE.

*Centaurea Jacea* L., var. *pratensis* Koch. Ditcham Park, Hants, August 25, 1914. A rather short small-flowered plant easily distinguished from the *C. decipens* and *C. nigra* forms growing with it. Heads sometimes rayed.—R. S. ADAMSON. Referred by Dr Thellung to *C. nigra* L. (pappus distinctus), sub-sp. *eumigra* Gugler. Folia insuete angusta.

?*Centaurea nigra* L., var. Rough pasture on clay, Ditcham Park, Hants, August 25, 1914. A very distinct small-headed form which seems near *C. Debeauxii* Gren. & Godr.—R. S. ADAMSON. "According to the somewhat artificial key of H. Saintange Savourié, this seems to

come under *C. consimilis* × *nemoralis*, though one of the parents may be *C. Debeauvii*, which occurs in S. England. My examples do not show fruit, which is useful in dealing with these *Centaurea* forms, but they agree with *C. consimilis* in the longly-ciliate, somewhat lax araneuse phyllaries, and with *C. nemoralis* in the deeper-coloured appendages and strongly inflated stem apex.—J. A. WHELDON. “Mr Adamson remarks:—‘A very distinct small-headed form, which seems like *C. Debeauvii* G. & G.’ I do not think this will do for *C. Debeauvii* G. & G., but I think it must be a *nigra* (including *obscura* Jord. and *nemoralis* Jord.) form, rather than coming under *pratensis* Thuill., in which group Rouy (*Fl. Fr.*) places *C. Debeauvii* as a ‘forme’ of *C. microptilon* Gren.”—C. E. SALMON. “*C. nigra* L. cf. sub-sp. *nemoralis* (Jord.) Gugler, acced. ad sub-sp. *Debeauvii* (G.G.) Gugler.” A. THELLUNG.

*Centaurea melitensis* L. Par, E. Cornwall, v.-c. 2, July 10, 1914. This plant was in good quantity at Par this year. Many plants were small, but there were three fine ones like good *Centaurea nigra*.—C. C. VIGURS. “Yes.”—A. THELLUNG.

*Centaurea*——? [Ref. No. 627.] On cotton-seed refuse with many other aliens, Hythe Quay, Colchester, N. Essex, v.-c. 19, June 16, 1914. Near *C. Calcitrapa* L., but flower bright yellow; no supplementary spines at base of large involueral spines; leaves with broader segments.—G. C. BROWN. “An early state of *C. Calcitrapa* L., is it not?”—E. S. MARSHALL. “This is *C. Calcitrapa* L.”—R. S. ADAMSON. “This is *C. pallescens* Del., var. *typica* Gugler and Thellung, forma *hyalolepis* Gugler. See Thellung *Adv. Fl. Montpellier* p. 546, 1912.”—A. THELLUNG.

*Centaurea aspera* L. Hayfield, Tottington, v.-c. 28, July 23, 1914.—F. ROBINSON. “Yes, var. *genuina* Willk.”—R. S. ADAMSON. “Recte.”—A. THELLUNG.

*Carthamus tinctorius* L. On rubbish by the canal, Litherland, S. Lancs, v.-c. 59, September 5, 1914.—J. A. WHELDON and W. G. TRAVIS. “Yes.”—A. THELLUNG.

*Picris Hieracoides* L., var. *gracilis* (Jord.). Letcombe Castle, Berks, August 1901. This variety is described by Rouy (*Fl. Fr.*, x., p. 23) as “plus grêle, pauciflore, à pubescence plus ténue, bien moins hispide; feuilles pubescentes, à peine rudes, plus brièvement dentées, calathides ± contractées à la maturité.” These characters we might assume to be caused by the habitat—dry chalk downs—on which it grew.—G. C. DRUCE.

*Crepis biennis* L. Edge of cultivated land, Walton, v.-c. 28, June 9, 1914.—F. ROBINSON. “Yes; not given for 28 in *Top. Bot.* This

plant seems to come under the var. *Bannatica* Rochel. Leaves unequally runcinate-pinnapartite or laciniate."—G. C. DRUCE.

*Crepis taraxacifolia* Thuill. [Ref. No. 46]. Cultivated land, light soil, Hargham, v.-c. 28, May 24, 1914.—F. ROBINSON.

*Hieracium Auricula* L. [Ref. No. 2832]. Originally from a pasture near Keevil, S. Wilts, v.-c. 8. Remote from houses, but only one patch seen. Cultivated at West Monkton, May 27, 1914. Styles yellow.—E. S. MARSHALL. "Correct."—E. F. LINTON.

*Hieracium cyathis* Ley. [Ref. No. 3975]. Cheddar Gorge, N. Somerset, v.-c. 6, May 30, 1914. Styles yellow. Leaves firm, rather glaucous, often blotched. Rev. Augustin Ley pointed out this to me several years ago as being his plant.—E. S. MARSHALL. "Exactly matches specimens in my herbarium gathered by the late Rev. A. Ley in the same Gorge on May 29, 1902."—J. CRYER. "This agrees with my plants from Cheddar, which were named *H. cyathis* for me by Rev. A. Ley."—E. F. LINTON.

?*Hieracium britannicum* F. J. Hanb. Limestone crags, the Red Screes, over Kirkstone Pass, Westmoreland, v.-c. 69, July 29, 1914.—C. WATERFALL. "No, young plants of *H. anglicum* Fr."—J. CRYER. "There is much resemblance to *H. britannicum* in these specimens, but the leaves of that species are rather strongly ciliate and toothed near the base, and the ligules are glabrous above. This plant is no doubt a variety of *H. anglicum* Fr., and probably var. *longibracteatum* F. J. Hanbury, but the specimens are undersized, having only one head each."—E. F. LINTON.

*Hieracium britannicum* F. J. Hanb. Ling Gill, Ribbleshead, alt. 1000 feet, v.-c. 64, July 25, 1914.—J. CRYER. "I cannot find anything better to put this to than *H. britannicum*, of which it may be a dry rock form. I have some exactly like it."—E. F. LINTON. "I think so."—E. S. MARSHALL.

*Hieracium sylvaticum* Fr. [Ref. No. 66]. Shady bank, West Runton, v.-c. 27, June 21, 1914.—F. ROBINSON. "No. 988, *L.C.*, ed. x., is *H. sylvaticum* Gouan. In my opinion, this is not *H. sylvaticum* Gouan, but *H. diaphanoides* Lindeb."—J. CRYER. "*H. sciaphilum* Uechtr., var. *transiens* Ley."—E. F. LINTON.

*Hieracium pellucidum* Laestad. Hackfall Woods, near Tanfield, v.-c. 64, June 7, 1913. This is a frequent hawkweed on the Carboniferous Limestone of Yorkshire, being found at Ingleton, Ribbleshead, Ling Gill, Hesledon Glen, Arncliffe, Kettlewell, Malham, Gordale, Grassington. The above is a new station for it in the North

Riding.—J. CRYER. "Rightly named."—E. F. LINTON. Also sent from Wyre Forest, Worcester, September 15, 1903. See *Report* 1903, p. 22.—Coll. AUGUSTIN LEY; comm. G. C. DRUCE.

*Hieracium lucidulum* Ley. Catterick Glen, W. Yorks, June 25, 1903.—Coll. AUGUSTIN LEY; comm. G. C. DRUCE. "*H. lucidulum* Ley = *H. pellucidum* Laest."—E. F. LINTON.

*Hieracium varicolor* Dahlst., stylose form. [Ref. No. 3309]. Originally from limestone, Allt nan Uamh, near Inchmadamph, W. Sutherland, v.-c. 108, July 1908. Flower garden, West Monkton, June 16, 1914. The original plants were so determined by the Rev. E. F. Linton. It comes true from seed. Styles greenish-yellow; heads epilose.—E. S. MARSHALL.

*Hieracium serratifrons* Almq., var. *caliginosum* Dahlst. [Ref. No. 3986]. Raised from seed collected by Mr W. A. Shoolbred near Inchmadamph, W. Sutherland, v.-c. 108, in 1908. Flower garden, West Monkton, June 8, 1914. Styles yellow; ligule-tips glabrous; heads very glandular, epilose; leaves dull, deepish green, white-dotted and glabrous above, very hairy beneath. Closely resembles plants so named from near Tongue and Kylesku, W. Sutherland.—E. S. MARSHALL. "I agree; the same form as we have so named for Mr Marshall from other places in Sutherlandshire."—E. F. LINTON.

*Hieracium sagittatum* Lindeb., var. *subhirtum* F. J. Hanb. Winch Bridge, altitude 900 feet, v.-c. 65, June 6, 1914. Teste Rev. E. F. Linton. A new county record, I believe.—J. CRYER. "Yes; heads larger than in the normal Scottish form."—E. S. MARSHALL. Also sent from High Force, Teesdale, altitude 1000 feet, v.-c. 65, June 6, 1914. Teste Rev. E. F. Linton.—J. CRYER.

*Hieracium*—? On the Carboniferous Limestone, Haweswater, Silverdale, v.-c. 60, May 30, 1914. The late Rev. A. Ley named similar specimens for me *H. cymbifolium* Purchas. The Rev. E. F. Linton writes on these specimens:—"I believe this is dwarf *H. expallidiforme* Dahlst., from limestone. It is worth cultivation to test proposed name."—J. CRYER. "Smaller than any specimen that I have of *H. sanguineum* Ley; but I think that it belongs here."—E. S. MARSHALL.

*Hieracium diaphanoides* Lindeb. Festiniog, Merioneth, July 14, 1903. See *Report* 1903, p. 22.—Coll. AUGUSTIN LEY; comm. G. C. DRUCE. "Of the three specimens on my sheet two are *H. vulgatum* Fr.; the third is probably a seedling of *H. diaphanoides*."—J. CRYER.

*Hieracium rigidum* Hartm., var. *Friesii* Hartm. Blaenau-Festiniog, Merioneth, July 14, 1903. See *Report* 1903, p. 23.—Coll. AUGUSTIN LEY; comm. G. C. DRUCE.

*Hieracium rigidum* Hartm., var.? Litton-dale, v.-c. 64, August 18, 1914. Mr Linton says:—"Var. *scabrescens* Dahlst. At least much the same as the Dent-dale specimens I so named for Prof. D. Oliver, and which W. R. LINTON accepts as that variety."—J. CRYER.

*Hieracium boreale* Fr. [Ref. No. 113]. Colwyn Bay, Denbigh, v.-c. 50, September 1, 1913.—W. C. BARTON. "Right."—E. F. LINTON. "Under the type, I believe; stem less hairy than usual."—E. S. MARSHALL.

*Hieracium boreale* Fr., var. *virgultorum* Jord. Wyre Forest, Worcester, September 15, 1903. See *Report* 1903, p. 23.—Coll. AUGUSTIN LEY; comm. G. C. DRUCE.

*Hieracium umbellatum* L., var. *pauciflorum* Hartm. [Ref. No. 49]. N. of Grand Havre, Guernsey, August 10, 1912. On specimens of this gathering submitted to him Mr Marshall reported:—"Practically identical with my specimen from Inveroran, Argyll (1893), named var. *pauciflorum* Hartm. by Elfstrand in F. J. Hanbury's herbarium." The Rev. E. F. Linton, who has seen the whole gathering writes:—"I agree with Mr Marshall's identification of your plant with his Inveroran specimen."—W. C. BARTON.

*Hieracium umbellatum* L.? var. *litorale* Fr. [Ref. No. 48]. Rocky coast, Cobo, Guernsey, August 16, 1912. Mr Marshall reports on a specimen of this gathering:—"I have exactly the same plant from rocky coast near Cobo Castle, and lane near Vale Castle (W. F. Miller, 1892). It may deserve a special name, but I know of none. Perhaps only a state of poor, rocky ground." Mr Linton, who has seen the whole gathering writes:—"Dr M. Elfstrand saw Cobo specimens of a dwarf *H. umbellatum* in F. J. Hanbury's herbarium, and was inclined to accept them as var. *litorale* Fr. Your plant looks to me more like var. *litorale* than the var. *littoreum* Arv.-Touv. of *Brit. Hieracia*."—W. C. BARTON.

*Hieracium umbellatum* L., var. *vincariifolium* Wallr. [Ref. No. 3997]. Locally plentiful among the coast sandhills, Ansdell, W. Lincs, v.-c. 60, August 10, 1914. Styles livid; leaves narrow, with revolute edges. Some of the stronger plants approach var. *coronopifolium* Fr.—E. S. MARSHALL. "I agree."—E. F. LINTON.

*Taraxacum erythrospermum* And. [Ref. No. 122]. Limestone rock, Bream Down, N. Somerset, v.-c. 6, April 26, 1914.—W. C. BARTON. "Yes; the exposed situation accounts for its compactness."—E. S. MARSHALL.

*Taraxacum spectabile* Dahlst. [Ref. No. 11419]. Fassaroe, Wicklow, April 1914. Growing in the demesne of our member, Mr R. M.

Barrington, with whom I gathered it last April. I suspect it to be allied to *spectabile*, and have sent it to Herr Dahlsted for his confirmation.—G. C. DRUCE. "I hardly think so. Leaf surfaces practically glabrous. No flowers on my two specimens; fruit not yet ripe. The habit is certainly not typical for *T. spectabile*. More like *T. palustre*."—E. S. MARSHALL.

*Campanula rotundifolia* L., var. *elongata* Hampe. Lough Gill, Sligo, August 17, 1913. Differs from my No. 51 sent last year from the same locality in the absence of the pubescence clothing the lower part of the stem. The height of these plants is remarkable (up to 80 cm.); the lower leaves about  $30 \times 7.5$  mm., while in the middle of the stem they are about  $60 \times 1.5$  mm.—W. C. BARTON. "This elongated state often occurs in Yorkshire. It is not worth a varietal name."—J. CRYER. "The variety is unknown to me; if this is it, I should suppose it to be a drawn-up 'state,' due to growing among rank vegetation."—E. S. MARSHALL. "Yes, but I am not certain whether it is more than a condition of growth."—G. C. DRUCE.

*Lobelia Dortmanna* L. Out of River Brathay, near Brathay Church, N. Lancs, v.-c. 69b, July 24, 1914.—C. WATERFALL.

*Calluna vulgaris* Hull, *forma*. [Ref. No. 629]. On soil which is inundated every winter. Tiptree Heath, N. Essex, v.-c. 19, August 13, 1914. Plants small, prostrate, cymes ascending; whole plant very pubescent, lighter green than type, which was in full bloom at time of gathering.—G. C. BROWN. "Obviously a 'state,' due to the situation."—E. S. MARSHALL.

*Daboecia cantabrica* R. & B. Lower slopes of Errisbeg, Roundstone, Galway W., August 13, 1913.—W. C. BARTON.

*Limonium vulgare* Mill, var. *pyramidale* Druce. Salt marsh, Keyhaven, S. Hants, August 1914.—J. COMBER. "Two specimens on my sheet. One is type; the other may pass as the 'variety.'"—C. E. SALMON. "My specimen is very weak, but may pass."—E. S. MARSHALL.

*Limonium binervosum* C. E. Salmon. Low place by sea, Hunstanton, v.-c. 28, July 31, 1914.—F. ROBINSON. "Yes, *Statice binervosa* G. E. Sm."—C. E. SALMON.

*Limonium recurvum* C. E. Salmon. Portland Island, v.-c. 9, July 14, 1914, with *L. binervosum*. Some of it was very much dwarfed. In fair quantity, and (I understand) a fairly safe situation. Some specimens of *Limonium* on the spot seemed to be rather off ordinary *binervosum* towards *recurvum*.—H. J. RIDDELSDELL. "Small and rather untypical, but I think correctly named."—C. E. SALMON.

*Primula elatior* Jacq. Wood, Great Sampford, N. Essex, v.-c. 19, April 10, 1914.—G. C. BROWN.

*Gentiana verna* L. Widdy Bank Fell, v.-c. 66, in abundance, altitude 1500 feet, May 1, 1914.—J. CRYER.

*Nymphodes peltatum* Rendle & Brit. Shallow water, Scoulton Mere, v.-c. 28, July 16, 1914.—F. ROBINSON. "The authority for *Nymphodes peltatum* is Otto Kuntze in his *Rev. Gen. Pl.*"—G. C. DRUCE.

*Amsinckia intermedia* F. & M. (*vide* Kew). [Ref. No. 128.] Mildenhall, W. Suffolk, v.-c. 26, June 6, 1913.—W. C. BARTON. "Probably, but the allied species are with difficulty distinguished."—A. THELLUNG.

× *Symphytum caeruleum* Petitmengin (*S. officinale* L. a. *ochroleucum* × *peregrinum*). See Bucknall's *Revision*, p. 550. Cultivated at Clifton, July, August, 1914.—J. W. WHITE.

*Symphytum grandiflorum* DC. (*S. ibericum* Stev.). Cultivated University Botanic Garden, Bristol, June, July 1914.—J. W. WHITE.

*Linaria Elatine* Mill. Damp places on edge of cultivated land, Watton, v.-c. 28, August 16, 1914.—F. ROBINSON.

*Veronica Anagallis* L. ? var. *montioides* (Boiss.) Hiern. Damp hollows in sand dunes, Freshfield, S. Lanes, v.-c. 59, July 19, 1914.—W. G. TRAVIS. "Probably so. I have gathered the same thing at Westenhanger, East Kent."—E. S. MARSHALL. "Yes, but is it more than a seedling state?"—G. C. DRUCE.

*Euphrasia nemorosa* Pers. (*vide* J. A. Wheldon). Steep hilly field on line side near Broadsands, Churston, S. Devon, v.-c. 3, July 7, 1913.—C. WATERFALL. "No; that is glabrous-leaved, whereas this is decidedly pilose. *E. curta* Wettst., var. *glabrescens* Wettst."—E. S. MARSHALL.

*Euphrasia curta* Wettst., b. *glabrescens* Wettst. (*vide* J. A. Wheldon). Open, bare hillside near Daddy Hole Plain, Torquay, S. Devon, v.-c. 3, June 16, 1913.—C. WATERFALL. "In bad condition, but correct."—E. S. MARSHALL.

*Euphrasia occidentalis* Wettst. ? Cliffs above Tilly Whim, Dorset, July 20, 1914.—A. B. JACKSON. "Yes; but plants not well selected."—E. S. MARSHALL.

*Euphrasia salisburgensis* Funck. Coast of Connemara, Ireland, August 1906.—J. W. WHITE. "Excellent specimens."—E. S. MARSHALL. "Yes, the most typical examples I have yet seen."—G. C. DRUCE.

*Euphrasia Kernerii* Wettst. Chalk hills near Reigate, Surrey, Sept. 7, 1913.—A. B. JACKSON, C. E. SALMON, J. FRASER. "Yes."—E. S. MARSHALL.

*Euphrasia foulaensis* Towns. (*vide* E. S. Marshall). [Ref. No. 266]. Short pasture at top of cliffs at seashore, Grim Ness, South Ronaldshay, Orkney, altitude 50 feet, July 15, 1914. Native. Leaves fleshy; cauline 2—6 toothed.—H. H. JOHNSTON.

*Euphrasia gracilis* Fr. [Ref. No. 118]. Marshy ground, Grande Mare, Guernsey, July 31, 1914. On specimens of this gathering submitted for comment Mr Marshall remarked:—"I think that this may be *E. gracilis* Fr., but am not sure. As a rule, that is a plant of rather dry ground." Mr Bucknall replied:—"E. *gracilis* Fr., I think." Most of the plants were growing in permanently water-logged soil, some in permanent water.—W. C. BARTON.

*Bartsia alpina* L. In abundance on Widdy Bank, Teesdale, v.-c. 66, altitude 1500 feet, June 6, 1914.—J. CRYER.

*Rhinanthus stenophyllus* Druce (*Alectorolophus stenophyllus* Sternneck). [Ref. No. 3934]. Meadows near Crianlarich, Mid Perth, v.-c. 88, July 18, 1914.—E. S. MARSHALL.

*Melampyrum pratense* L., agg., var. [Ref. No. 11149]. Millook, Cornwall, June 1914.—G. G. DRUCE. "My largest example is var. *latifolium* Schreb. & Mart., which Dr Moss regards as the type of the species, or near it. The smaller pieces only differ in being depauperate."—E. S. MARSHALL.

*Orobanche major* L. Near Cheddington, Bucks, August 1904.—G. C. DRUCE.

*Utricularia major* Schmidel. Pools at Restennet, Forfar, v.-c. 90, August 4, 1914.—R. & M. CORSTORPINE. "No note as to the veining of the flowers, which is conclusive as to its being *major*. I suppose it must be so named, but the bladders are large, though not so large as in the *F. gigantea* Prah. I think this is a new record for Forfar county 90, as Mr Marshall's specimens thence were rather doubtful."—A. BENNETT. "Yes, excellent examples showing the 'winter buds' as well as the flowers."—G. C. DRUCE.

*Mentha longifolia* Huds., b. *nemorosa* (Willd). [Ref. No. 85].  
By ditch, Carbrooke Fen, v.-c. 28, August 20, 1914.—F. ROBINSON.  
"So I should name it. Leaves rather short."—E. S. MARSHALL.

*Mentha rotundifolia* × *longifolia* = *villosa* Huds. Roadside near  
Dilty Moss, Forfar, v.-c. 90, September 7, 1914.—R. & M. CORSTORPHINE.  
"I believe this is correctly named."—C. E. SALMON. "Seems  
rather a good intermediate."—E. S. MARSHALL.

*Mentha citrata* Ehrh. Origin Northaw, Herts (H. Peirson), Hort.  
Reigate, Surrey, September 1914. This is the plant of Pryor's *Fl.*  
*Herts* (1887), p. 336, where it stands as a variety of *M. hirsuta*.  
These specimens show the globular heads of *M. aquatica* (*hirsuta*) and,  
whilst not absolutely glabrous, the whole plant is more so than in Mr  
White's "*citrata*" from Priddy Nine Barrows (*Rep.* for 1908, p. 393,  
and *Wats. B.E.C. Rep.* for 1909-10, p. 250). It will be noted, too,  
that the leaves are broader and more cordate than in Mr White's  
plant—all points in favour of *citrata*. Ehrhart's original description  
is extremely terse—"Folia ovata, petiolata, serrata, glabra. Capitula  
obtusata. Stamina corolla breviora." *Beitr.* vii. (1792), p. 150.—C. E.  
SALMON.

*Mentha hirsuta* L., var. *sub-glabra* Baker. Crombie Den, near  
Arbroath, v.-c. 90, September 20, 1914.—R. & M. CORSTORPHINE.  
"Judging by examples named by Mr Baker I believe he would pass  
this as his variety."—C. E. SALMON. "Probably. I have not seen  
Baker's plant. A form, apparently referable to this, found in 1906 at  
Braemar by W. A. Shoobred and myself, had the sweet scent of  
*M. citrata*."—E. S. MARSHALL. "Yes, but not extreme."—G. C.  
DRUCE.

*Mentha Pulegium* L., var. *erecta* Syme. Marshy ground, Limps-  
field Common, Surrey, August 23, 1914.—A. B. JACKSON and J.  
FRASER. "This agrees much better with Rouy's description of *β.*  
*vulgaris* Briquet than with var. *erecta* Wirtgen, whose name antedates  
Syme's."—E. S. MARSHALL. "Yes, but Briquet in his *Labiées*, p. 93,  
gives the authority for *erecta* as Wirtgen."—G. C. DRUCE.

*Thymus Serpyllum* L., var. *Linneanus* G. & G. [Ref. No. 126].  
Edge of road, Albecq, Guernsey, July 31, 1914.—W. C. BARTON.  
"This, I think, is our assumed type."—E. S. MARSHALL.

*Stachys palustris* × *sylvatica*. [Ref. No. 1238]. Green Lane,  
Merton parish, Surrey, July 18, 1914.—C. E. BRITTON. "Yes; con-  
siderably nearer to *S. palustris*."—E. S. MARSHALL. "Yes; and very  
like the figure of Smith's *ambigua* in *E.B.*, t. 2089."—C. E. SALMON.  
Also from bank of stream, the Hall Dell, Melmerby, Cumberland,

v.-c. 70, July 31, 1914.—Coll. Rev. W. W. MASON; comm. C. WATERFALL. Also from ditch near Duncan's Cottages, Billingshurst, Sussex, July 14, 1914.—A. WEBSTER. "I believe so; but far closer to *S. palustris*."—E. S. MARSHALL. Also from the banks of the Leeds and Liverpool Canal, Shipley, v.-c. 64—with both parents, August 4, 1914.—J. CRYER. "*S. palustris* × *sylvatica*, much on the *palustris* side."—E. S. MARSHALL. "The specimens from Messrs Britton, Waterfall, Webster, and Cryer, all come under this hybrid."—G. C. DRUCE.

*Galeopsis angustifolia* Ehrh., var. *canescens* Schultes. [Ref. No. 114]. Avebury Down, N. Wilts, v.-c. 7, August 5, 1914.—W. C. BARTON. "Yes; if Koch's description of this variety is correct. He states that *canescens* has short, dense, patent hairs; *angustifolia*, 'pili omnes adpressi.' The former is surely our more common plant by far."—C. E. SALMON.

*Leonurus Cardiaca* L. Open wood and roadside, Hargham, v.-c. 28, June 29, 1914.—F. ROBINSON.

*Lamium maculatum* L. Hedgebank, roadside, Ovington, v.-c. 28, April 26, 1914.—F. ROBINSON.

*Lamium purpureum* L. Cultivated Dalton, v.-c. 69b, August 7, 1914. See *Report* 1913, p. 491. The following characters are maintained:—The long cotyledons; the variably fissile lower lip of the corolla; the asymmetrical sub-rhomboidal leaves; the absence of any cordate base; the shallow, very irregular toothing; the "dappling" of the leaves; the almost obsolete rugosity; the thin texture; the short leaf-stalks—half inch at most. The plant is a very "shy" seeder; nearly all the seeds produced germinate.—D. LUMB. "A very curious form, worth further study and perhaps a varietal name."—G. C. DRUCE.

*Lamium purpureum* L., ?var. *decipiens* auct. Denham, Bucks, June 1902.—G. C. DRUCE. "Apparently a true variety of *L. purpureum* rather than its cross with *L. hybridum*."—E. S. MARSHALL.

*Ballota ruderalis* Koch. Llamwarne, Herefordshire, July 18, 1914.—A. WEBSTER. "No, the teeth of the calyx in *ruderalis* should be from 2—4 mm. long; these are not 2 mm. The calyx in *ruderalis* is more hairy. This is *B. nigra*."—G. C. DRUCE.

*Teucrium Scordium* L., var. Braunton Burrows, N. Devon, August 1896.—G. C. DRUCE.

*Plantago ceratophylla* Hoffmg. and Link. Seedlings. Cultivated Walton, S. Lancs, July-September 1914.—J. A. WHELDON.

*Plantago Coronopus* L., var. [Ref. No. 4050]. Sandstone cliffs, Sidmouth, S. Devon, v.-c. 3, June 8, 1914. Very large (all my gathering came from two plants); biennial.—E. S. MARSHALL.

*Plantago Coronopus* L., form or var. [Ref. No. 4049]. Sandy ground near the coast, Berrow, N. Somerset, v.-c. 6, June 4, 1914. Apparently biennial; closely appressed to the ground. May be only a peculiar state of the type.—E. S. MARSHALL. "*P. Coronopus* L., var. *transiens* Beguinot."—R. M. CARDEW and E. G. BAKER.

*Plantago Lagopus* L. [Ref. No. 717.] On cotton-seed refuse, Hythe Quay, Colchester, N. Essex, v.-c. 19, June 1914. Apparently referable to this, though the dark, villous part of the dorsal portion of the sepal is narrower than in that of an old specimen from the Isle of Elba in my herbarium.—G. C. BROWN. "Correct."—E. G. BAKER. "Yes, forma *glabris* (corollæ lobis extus glabris nec pilosis)." —A. THELLUNG.

*Plantago major* L., var. *intermedia* (Gilib.) Barnards Green, near Malvern, Worcester, v.-c. 37, August 17, 1914.—Coll. R. F. TOWNDROW; comm. S. H. BICKHAM. "*Plantago major* L., var. *intermedia* Syme."—R. M. CARDEW and E. G. BAKER. "My single specimen differs from the description of *P. intermedia* in Williams *Prodr. Fl. Brit.*, pp. 354-5. Not very hairy; leaves not coarsely toothed towards the base, &c. But it comes near what is usually so named with us.—E. S. MARSHALL. Also sent from Sands of Barry, v.-c. 90, Aug. 10, 1914.—R. & M. CORSTORPHINE. "The *P. intermedia* of Gilibert is an entirely different plant. It is figured and described by this author in his *Histoire des Plantes d'Europe* in 1806. His figure shows a plant with sessile strongly serrate leaves, reaching 2 in. long, and 1 inch broad. The scapes are hairy, and the spikes dense and short. The whole plant rather suggests a form of *P. media* L."—R. M. CARDEW and E. G. BAKER.

*Plantago major* L., var. *minima* DC. Malvern Common, Worcester, v.-c. 37, August 17, 1914.—Coll. R. F. TOWNDROW; comm. S. H. BICKHAM. "Does not agree at all well with the description in DC. *Prodromus*, xiii. a., p. 695:—'parvula pollicaris, foliis ovatis integris glabris in petiolum brevem attenuatis, pedunculis folia subaequantibus erectis vel ascendentibus, spicis ovatis vel ovato-capitalis.'" The sepals are rounded-obtuse; otherwise it is much like the plant sent as var. *intermedia* on a reduced scale."—E. S. MARSHALL.

*Illecebrum verticillatum* L. Pine Wood near Wellington College, June 6, 1914.—A. WEBSTER.

*Herniaria ciliata* Bab. Lizard, W. Cornwall, v.-c. 1, August 20, 1914.—Coll. E. THURSTON; comm. C. C. VIGURS.

*Chenopodium murale* L. Trent Meadows, Nottingham, October 7, 1914. This species was growing in fair abundance on waste ground adjacent to the site of the abortive Nottingham Exhibition of 1913. In the same field *Crocus nudiflorus* is abundant, and appears to be on the increase, owing to the erecting of palings for the above purpose around the station. With *C. murale* grew *C. opulifolium*, *C. olidum*, *C. rubrum*, *C. album*, vars., etc., and many interesting casuals, such as *Glaucium luteum*, an unexpected alien inland.—A. R. HORWOOD. "Yes."—G. C. DRUCE. "My specimen is abnormal (probably injured); it may be the sub-var. *microphyllum* Coss. and Germ. = var. *microphyllum* Gürcke, being very small-leaved, but its condition does not admit of certainty."—E. S. MARSHALL. Also sent from Newquay, W. Cornwall, v.-c. 1, December 1913.—C. C. VIGURS. "Yes."—C. E. SALMON, E. S. MARSHALL, and G. C. DRUCE.

*Chenopodium ficifolium* Sm. Oakfield, Hitchin, Herts, v.-c. 20. With *C. album* and perhaps hybridising, September 9; fruit October 17, 1914.—J. E. LITTLE. "Surely not *C. ficifolium*, but a form of *C. album* L."—J. CRYER. "Yes."—E. S. MARSHALL.

*Chenopodium Vulvaria* L. Seedlings. Albecq, Guernsey, July 31, 1914.—W. C. BARTON.

*Chenopodium hircinum* Schrad., var. *subtrilobum* Issler. Wool waste heap, Selkirk, v.-c. 79, October 1913. Teste Dr Murr.—I. M. HAYWARD.

*Beta trigyna* Waldst and Kit. Waste ground, St Philips, Bristol, June 20, 1914.—J. W. WHITE. "Yes."—G. C. DRUCE.

*Salicornia ramosissima* Woods. Mud flats, Montrose Basin, v.-c. 90, September 20, 1914.—R. & M. CORSTORPHINE. "I would suggest that this is a small boreal form of *S. herbacea* L., forma *patula* Moss."—E. S. MARSHALL.

*Salicornia ramosissima* Woods, forma. Mud flats, Montrose Basin, v.-c. 90, September 20, 1914.—R. & M. CORSTORPHINE. "Surely not; spikes too obtuse. I think this to be *S. gracillima* Moss; new for Scotland."—E. S. MARSHALL.

*Salicornia dolichostachya* Moss. Mud flats, Montrose Basin, v.-c. 90, September 20, 1914.—R. & M. CORSTORPHINE. "Yes, an interesting extension of its range in Great Britain. Sent to me fresh by the collectors last autumn from the same locality. It occurs in Denmark."—C. E. MOSS. "Just like plants gathered (in 1914) near Emsworth, Hants, which Dr Moss confirmed as being this species. Evidently closely allied to *S. stricta*."—C. E. SALMON. "Very characteristic;

new for Scotland, I think. For years a specimen from Sandwich flats, E. Kent, collected by Mr G. Dowker, perplexed me; it is a small form of this species."—E. S. MARSHALL.

*Salicornia prostrata* Pallas, var. *Smithiana* Moss and Salisbury. Mud flats, Montrose Basin, v.-c. 90, September 20, 1914.—R. & M. CORSTORPHINE. "I think so."—E. S. MARSHALL.

*Polygonum lapathifolium* L., var. Cornfield, Wigginton Heath, Oxon, July 29, 1914. Glands on perianth few, but the nut is that of this species. The variety deserves a name if a name is ever to be given for hairy leaves. Flowers darker than usual.—H. J. RIDDELSDELL. "Too slender, I believe, for any *lapathifolium* form; it may be *P. nodosum* Pers., forma *salicifolium* Moss (*P. incanum* Willd.), though the stems are unspotted, and not swollen at the nodes; or else a hybrid of that with *P. Persicaria*, to which it bears much resemblance. The peduncle-glands are numerous, though shortly stalked."—E. S. MARSHALL. "Under leaves with whitish down approaching var. *incanum*."—G. C. DRUCE.

*Polygonum* sp. In an elevated sandy cornfield, c. 650 feet, on Wigginton Heath, Oxon, August 17, 1914. Characterised by the remarkably narrow acute leaves and silvery ocreæ.—H. J. RIDDELSDELL. "*P. heterophyllum* Lindm., *sensu lato*—probably a form of *P. ruriavagum* Jord."—G. C. DRUCE.

*Polygonum*—? Waste, stony ground, courtyard of Imperial Institute, September 19, 1914.—A. B. JACKSON.

*Polygonum aviculare* L., var. *ruriavagum* Jord. Weed in field, Grey Abbey, County Down, September 27, 1914.—C. H. WADDELL. "*P. heterophyllum* Lindm. forma; not, I think, *ruriavagum* Jord."—G. C. DRUCE.

*Rumex salicifolius* Wien. Walton, S. Lanes, August 20, 1914.—J. A. WHELDON.

*Ulmus scabra* Mill. Millwood, Dalton, v.-c. 69b, September 4, 1914. Two year old plants.—D. LUMB.

*Ulmus stricta* Lindley. The Cornish Elm. Penpoll Quarry, Cranstock, Newquay, W. Cornwall, March 15, 1914. These flowering specimens from the same tree from which foliage was distributed in 1911 may be acceptable. This tree fell into a horizontal position some years ago, so that one can get at the top branches fairly easily. I hope to get fruit from it some year. The Cornish Elm does not fruit as freely as the Dutch, and even when it does the fruit is not easy to get, unless one is a very expert tree climber. I should be inclined to object to

the term "pyramidal" applied to the outline of the Cornish Elm in the *Cambridge Flora*. I should say "cylindrical" would be a better term, though that is hardly correct, the longest branches being usually just below the top, and all the lower branches short and scattered. This is probably the reason why rooks prefer this tree to any other for nesting in.—C. C. VIGURS.

*Ulmus glabra* Huds. = *U. montana* Stokes. Trevowah, Crantock, Newquay, W. Cornwall, middle of May 1914. These specimens in good fruit are from one of the very few (Davey's *Flora* notwithstanding) trees of this species in the county authoritatively named since the elms became better understood than they used to be. I think this species used to be confused with *U. hollandica*. This particular tree, named by Dr Moss, fruits well.—C. C. VIGURS. "The petioles are rather long for *U. montana*. One would like to see examples with mature leaves."—A. B. JACKSON. "Looks right."—E. S. MARSHALL.

*Alnus rotundifolia* Mill, var. *incisa*. A fine tree, 40 feet high, planted in the Rectory Close, Wigginton, Oxon, September 1, 1914. Is this form of the alder native to the British Isles? Babiington, ed. 9, gives it for Wigtonshire apparently as a native.—H. J. RIDDELSDELL. "The inflorescence is that of *A. glutinosa* Gaertn., var. *microcarpa* Rouy."—E. S. MARSHALL. "This is not the var. *incisa*, which has the leaves small and deeply incised, resembling those of the common hawthorn, but is the cut-leaved alder, *A. glutinosa* var. *laciniata* Willdenow *Berlin Baumz.* 44, 1796, which is frequent in cultivation and often attains a large size, as at Syon House and Woburn. According to Duhamel it occurs wild in the north of France, particularly in Normandy, and in the woods of Montmorency near Paris."—A. B. JACKSON. "In Oxfordshire var. *laciniata* Willd. only occurs as a planted tree, and there are fine examples in "The Parks" at Oxford. Wm. Cobbett is said to have planted a tree at Wolvercote. See *Report* 1909, p. 473. Dr Balfour was the authority for its occurrence in Wigtonshire (? native). Our late member, T. A. Stewart, found a specimen on the Black Mountain, Belfast, but he does not mention it in his *Flora*. It occurs (? planted) at Lakenham Bridge, and is said to be native in Northern France."—G. C. DRUCE.

*Salix triandra* L. [Ref. No. 225]. Hedge, Earls Colne, N. Essex, v.-c. 19; flowers, April 30; leaves, August 23, 1914. Teste G. C. Druce.—G. C. BROWN. "Rightly named."—E. F. LINTON. "Type (*genuina* Syme) I believe."—E. S. MARSHALL.

*Salix triandra* L., ♂ *forma*. [Ref. No. 229]. "The Moors," Alphamstone, N. Essex, v.-c. 19, May 1914. See *B.E.C. Report* 1913, p. 496. I send a supply of male flowers to supplement leaf-specimens

(September 21, 1913) sent last year from the same bush. The bush flowered profusely this year.—G. C. BROWN. "No mature leaves; but it looks like var. *Hoffmanniana* Bab."—E. S. MARSHALL.

*Salix purpurea* L., ♂. [Ref. No. 227.] Earls Colne, N. Essex, v.-c. 19; flowers, April 30; leaves, August 23, 1914.—G. C. BROWN. "Correct."—E. F. LINTON. "Seems to come under the type (*vera* Ritschl)."—E. S. MARSHALL.

*Salix Forbyana* Sm. Lowland near Brent Knoll, N. Somerset, April and August 1902.—J. W. WHITE. "Rightly named."—E. F. LINTON. "A broad-leaved *S. purpurea* × *viminalis*, which I think answers to this name."—E. S. MARSHALL.

*Salix aurita* aut *cinerea* × *viminalis*? = *S. ferruginea* G. Anderson? Meadows near Long Ashton, N. Somerset, v.-c. 6, April, May, and August 1913.—J. W. WHITE. "The male specimen is *S. cinerea* × *viminalis*; the female probably *S. aurita* × *viminalis*. The foliage looks rather as if it belonged to the latter, but is wanting in decisive features, and there is nothing to show whether it is from the male or the female bush, or from either. I have remarked before on the objectionable practice of sending male and female specimens on the same sheet, unless, of course, there is no doubt that both belong to the same species. In this case the result of the mixture gives a very unsatisfactory result."—E. F. LINTON. "This I consider to be *S. cinerea* × *viminalis*."—E. S. MARSHALL.

*Salix phylicifolia* (L.) Sm. [Ref. No. 1822.] (Name confirmed by E. F. Linton). Grassy banks at burnside, altitude 50 feet, Burn of Stennadale, Firth, Mainland, Orkney; flowers, June 3; leaves, August 5, 1914. Native, common. A straggling shrub with sub-erect or erect stems, 1—2 feet high.—H. H. JOHNSTON.

*Ceratophyllum submersum* L. Pond, Castle Morton, Worcester, v.-c. 37, August 27, 1914. I understand that Dr C. E. Moss prefers to call the *C. submersum* var. *apiculatum* of Dalla Torre and Sarntheim. Coll. R. F. TOWNDROW; comm. S. H. BICKHAM. "This plant is, in my opinion, *C. demersum*, var. *apiculatum* = *C. apiculatum* Cham. I have never gathered *C. submersum* in any inland county, and have only seen British specimens from southern England—Somerset to Norfolk. The var. *apiculatum* is intermediate between *C. demersum* and *C. submersum*, and it is somewhat arbitrary to refer it to one of the species rather than the other, but I prefer to put it to *C. demersum*, simply because this plan enables one to determine the two species in absence of ripe fruit. I agree that in fruit characters the variety is not far from *C. submersum*, and I should not complain if the two species were reduced to one."—C. E. MOSS.

*Ceratophyllum demersum* L. Pond, Little Malvern, Worcester, v.-c. 37, August 25, 1914. Teste C. E. MOSS—*i.e.*, specimens from the same pond were passed. Dr MOSS has not seen these individual examples.—Coll. R. F. TOWNDROW; comm. S. H. BICKHAM.

*Epipactis palustris* Crantz, var. *ericetorum* Asch. & Grabn. Sandhills, Ainsdale, S. Lancs., v.-c. 59, July 5, 1914.—W. G. TRAVIS. "In my opinion, merely a state due to environment."—J. CRYER. "Approaching *Helleborine palustris*, var. *ericetorum* (A. & G.) Druce, but more luxuriant than the specimen I gathered there in 1911, and with narrower leaves."—G. C. DRUCE.

*Orchis incarnata* L. Damp flats, Sands of Barry, v.-c. 90, June 13, 1914.—R. & M. CORSTORPHINE. "Yes, a small form."—E. S. MARSHALL.

*Orchis praetermissa* Druce. [Ref. No. 643.] Meadows, Henny, N. Essex, v.-c. 19, July 2, 1914. Flowers varying from pale rose-purple to deep purple; lip almost flat, varying somewhat in markings and shape, but always as broad as long. Many of the bracts are coloured. Despite every care, I was unable to preserve the colour of the flowers.—G. C. BROWN. "In Gibbons Brook, Kent, at Cray, in Mid-West Yorks, and near Silverdale, in Lancashire, I have found every shade of colour in the flowers of different specimens from light flesh colour to 'dark crimson purple,' and apart from the shade of colour I find no distinguishing character whereby they can be separated. In these three stations the plants with different shades of colour bloom at the same time. The sides of the lip in every instance were reflexed, and so they are in this specimen. I should call it *Orchis incarnata* L."—J. CRYER. "As I have not yet seen the description, these specimens are very welcome. For many years I have believed that our *incarnata* included two or more species. As far as one can judge from dried material, this is what Max Schulze (*Orchidaceen Deutschlands und der Schweiz*, tab. 19) figures as *O. incarnata* L.—not the Linnean plant. I have found it in several southern and eastern counties."—E. S. MARSHALL. "The middle lobe of labellum is longer than in the type, and suggests the presence of *maculata*. I should like to see it in the fresh state."—G. C. DRUCE.

*Aceras anthropophora* Br. Calcareous pastures, near Barnack, Northants, v.-c. 32, May 30, 1914. The stems of most of the specimens gathered had been withered by very late frosts. It is rather curious that the orchis should have been surprisingly abundant this year.—G. CHESTER. Also from chalk downs, Reigate, Surrey, May 31, 1913.—C. E. SALMON.

*Ophrys apifera* Huds. Clay soil, railway cutting, Saham Toney, v.-c. 28, June 24, 1914.—F. ROBINSON.

*Habenaria virescens* Druce. Damp wood, Griston, v.-c. 28, June 17, 1914.—F. ROBINSON.

*Leucojum vernum* L. [Ref. No. 4056]. Discovered by Miss M. A. Hellard, between Bishop's Lydeard and Williton, S. Somerset, v.-c. 5, February 24, 1914. A very welcome novelty for this county. It grows in good quantity over a limited area.—E. S. MARSHALL.

*Asparagus officinalis* L., *a. attilis* L. Sandhills, Freshfield, v.-c. 59, July 1914.—W. G. TRAVIS.

*Allium sphaerocephalum* L. St Vincent's Rocks and Durdham Down, Bristol, July 27 and 31, 1911. The larger-headed specimens are from the former locality, and those with smaller heads from the latter. The plant is too scarce for any extensive gathering, and no bulbs have been taken for the Club. The few now sent had been pulled up and thrown aside by scrambling boys, who, in attempting to gather the flowers, had uprooted the plant from the loose thin soil on the rocks.—J. W. WHITE.

*Allium sibiricum* L. Predannock Downs, Lizard, West Cornwall, v.-c. 1, June 11, 1914.—Coll. E. THURSTON; comm. C. C. VIGUES.

*Allium Schænoprasum* L. "Chivey Syke," v.-c. 69b, June 8, 1914. I send these few flowers, without bulbs, to confirm Miss Hodgson's old record. A vandal, at one visit, could easily make the plant non-existent here.—D. LUMB.

*Allium sativum* L., var. *Ophioscorodon* Doell. [Ref. No. 4905]. Ex Mull of Galloway, July 1912.—G. C. DRUCE.

*Fritillaria Meleagris* L. Damp shady pasture, Ovington, v.-c. 28, April 27, 1914.—F. ROBINSON. "Not given for West Norfolk in *Top. Bot.*"—G. C. DRUCE.

*Paris quadrifolia* L. Damp wood, Seamere Wood, Hingham, v.-c. 28, March 21, 1914.—F. ROBINSON.

*Juncus maritimus* Lam., var. *atlanticus* mihi. Salt-marsh, St Mary's, Scilly, September 5, 1914. By the kindness of Mrs Stideford of "Lunnon" I am enabled to distribute another parcel of this interesting rush. My correspondent secured some good stems before the marsh was mown, but has cut them shorter than is perhaps desirable. In my note on this plant (*Journ. Bot.* January 1914, p. 19) I proposed for it the varietal name *atlanticus*, having concluded that the allied form *J. rigidus* Desv. (*Rouy Fl. de France*), described as "*forte, rigide*," could not be identical. That description indeed

seems to fit the type *maritimus* of this country rather than the variation under notice, which has a rather weak slender stem from four to five feet high. Still, as Dr Moss has suggested, it will be well to compare this plant with specimens of *J. rigidus* in the Rouy Herbarium at Paris when an opportunity offers; and until that can be done the name *atlanticus* should be regarded as provisional. Examples in some degree approaching the Scillonian form have been lately forwarded from Poole Harbour, Dorset, by my friend Mrs E. P. Sandwith. The following brief description may suffice to define this variety:—*Culmo subtenui, elato, ad 10—15 dcm. producto. Anthela magna (2½—4 dcm. longa) diffusa, abunde decomposita, bracteam floralem inferiorem multo superante. Cætera ut typi.* With reference to the comments of Professor Lindman and Mr Adamson in the *Report* for 1913, p. 499, I would say that no botanist who had seen it growing in masses over a large area could possibly suppose this plant to be a monstrosity; and would ask how any state of luxuriance could so shorten the lower bract, not merely in relation to its own panicle, but in relation to that of a non-luxuriant type.—J. W. WHITE. “I find in my herbarium a specimen of this interesting variety, collected by Mr A. Somerville, in July 1890, in the same station (Mr White confirms the name). This shows that the variation is not merely a ‘state,’ ‘here to-day, gone to-morrow,’ which is interesting in the face of some of the comments on p. 499, *B.E.C. Rep.* 1913.”—C. E. SALMON. “It certainly is a peculiar form. Only one variety is given by Ascherson and Graebner in their *Syn. Flor. Mitt.-Eur.* 1904, p. 456, and that is a very local plant.”—A. BENNETT. “A well-marked variety, in my opinion; contrasting greatly with a slender, narrow-panicked plant found on the Lancashire coast-sands.”—E. S. MARSHALL.

*Juncus filiformis* L. Derwentwater, Cumberland, August 1902.—G. C. DRUCE.

*Juncus bulbosus* L., var. *uliginosus* (Fr.) Druce. [Ref. No. 103.] Wet lane, Petit Bo, Guernsey, August 6, 1912.—W. C. BARTON. “So I should name it.”—C. E. SALMON. “This is not var. *uliginosus*, which is a procumbent form seldom “bulbous” and with three stamens. The present plant has six stamens with the anthers only about one-third the length of the filaments. It would apparently come under the var. *Kochii* = *J. nigritellus* Koch non D. Don. This is generally described as having the capsule equal or shorter than the perianth, not much longer, as in the present case. I know this long-pointed form from many woodland districts. On further examination of the plants sent by Mr Barton last year from Roundstone, Co. Galway, it would appear they are the same as this, and not *J. bulbosus* f. *uliginosus*, as previously reported.”—R. S. ADAMSON. “I look upon this simply as a viviparous condition, not a true variety.”—E. S. MARSHALL.

*Juncus tenuis* Willd. Roadsides, Crianlarich, Mid Perth, v.-c. 88, July 15, 1914.—E. S. MARSHALL.

*Juncus bufonius* L., var.? Sandy shore, at high watermark, Ballywalter, Co. Down, August 1914. These plants are frequently overflowed by the tide. I suppose the sepals are too sharp for var. *ranarius*.—C. H. WADDELL. "This is apparently the plant described as var. *ranarius* = *J. ranarius*, but the original description of *J. ranarius* Song. et Perrier says that the perianth is shorter than the capsule and the inner obtuse, while in this plant the perianth segments are all acute and exceed the capsule. A specimen in Camb. Univ. Herbarium, issued by Perrier, of *J. ranarius* is a very different plant, that so far I have not seen in this country. The present plant would come under *J. ranarius* sensu Asch. and Graebner *Syn. Mitt. Fl. ii.*, 2, 1904, p. 432, or *J. insularis* Viv. in Rouy *Fl. Fr. xiii.*, 1912, p. 252, which Buchenau treats as the same as *J. bufonius* var. *fasciculatus* Koch. The oldest varietal name would seem to be var. *congestus* Wahlb. (1820), but I have not seen the description."—R. S. ADAMSON. "I have just the same thing from Starcross, S. Devon (October 1888); and New Romney, E. Kent (July 1891). Much paler than var. *fasciculatus* Koch, the individual flowers being as a rule fully twice as long. My only sheet labelled *J. ranarius* Nees, from Southport, S. Lancs, is too dwarf and poor to give much help, but I suspect that all these gatherings may belong to it."—E. S. MARSHALL. "This is what I called *fasciculatus*."—G. C. DRUCE.

*Juncoides Fosteri* Sm. Thicket near Colwall, Herefordshire, v.-c. 36, May 9, 1914.—C. WATERFALL. "Right."—E. S. MARSHALL. Also from Harefield, Middlesex, v.-c. 21, May 4, 1913.—W. C. BARTON.

*Luzula multiflora* DC., var. *congesta* (Lej.). Tiptree Heath, N. Essex, v.-c. 19, May 5, 1914.—G. C. BROWN.

*Sparganium neglectum* Beeby. By the Dane Stream, Milford-on-Sea, S. Hants, v.-c. 11, August 1914.—J. COMBER. "Probably correct, but less 'gradually attenuated' than in the original specimens of Mr Beeby."—A. BENNETT. "The specimen before me is useless, having no good fruit; I received a better one through the Watson Exchange Club. From the very numerous fruits, many of them somewhat angular (or 'shouldered') at the broadish base, I would name it *S. erectum*, var. *microcarpum*."—E. S. MARSHALL.

*Sparganium minimum* Fr. Borough Fen, Northants, August 1910. A new county record. Here over a small area it was plentiful.—G. C. DRUCE.

*Damascium Alisma* Mill. Hook, N. Hants, August 1910.—  
G. C. DRUCE.

*Scheuchzeria palustris* L. [Ref. No. 3941]. Bog near Rannoch Station, Mid Perth, v.-c. 88, July 17, 1914; fruiting freely. I afterwards found it in another swamp not far off, in leaf only.—E. S. MARSHALL.

*Potamogeton Zizii* Koch. Crooked drain, near Ely, Cambridge, June 2, 1884.—H. and J. GROVES. "Is *P. praelongus* Wulfen."—A. BENNETT.

*Potamogeton decipiens* Nolte. Canal, Market Harboro, Leicestershire, v.-c. 55, July 19, 1914.—G. CHESTER. "Yes, very fair example of Nolte's plant."—A. BENNETT.

× *Potamogeton Lintoni* Fryer. Canal, Renishaw, Derby, October 1911. A supposed hybrid of *crispus* and *Friesii*. It grew in considerable quantity and with some amount of variability in the Renishaw Canal, to which place I went in order to show Dr Glück × *Apium Moorei* in situ.—G. C. DRUCE. "Yes. *P. crispus* × *Friesii* Fryer. This was gathered near Sheen in Surrey in 1912, C. E. Salmon sp."—A. BENNETT.

*Potamogeton acutifolius* Link. Water course near Wareham, Dorset, June 30, 1914.—J. W. WHITE. "Yes; the smaller form of the species."—A. BENNETT. "Yes, from a well-known locality, in which I have gathered it."—E. S. MARSHALL. Also from deep pool, Staines, Middlesex, July 19, 1879, and July 29, 1882.—H. GROVES.

*Potamogeton trichoides* Cham. and Schlecht. Ditch near Horsey, E. Norfolk, July 10, 1912.—J. GROVES. "I suppose must be so named. It differs from the typical specimens in the Berlin Herbarium by the leaves being mostly three-veined and the fruit not so broad, and the projecting pit being in the circumference of the fruit, not outside it. It differs from the usual British plant by not being tuberculate on the dorsal margin, this being the var. *Trimmeri* of Dr Caspary."—A. BENNETT. "Evidently later-flowering than most of our species; no fruit present the second week in July."—E. S. MARSHALL.

*Potamogeton marinus* L. Coldisham Loch, Berwick, August 1906. In great quantity, preferring the shelter of *Castalia*. I use the older and, I believe, the correct Linnean name in preference to that of Nolte.—G. C. DRUCE. "Yes, characteristic examples of Nolte's *P. filiformis*."—A. BENNETT.

*Naias flexilis* R. & S. Esthwaite Water, v.-c. 69b, August 14, 1914. The dominant species in a remarkable association (see *Hydrilla verticillata*). Although the species fruits freely it is so abundant that it must have been established here for some considerable time.—W. H. PEARSALL. "An excellent addition to the English flora. Mr Pearsall sent me the first specimens found, asking if it was not *Naias*, and with these specimens a scrap of what I at once saw could only be *Hydrilla verticillata* Caspary, which was an addition to the flora of the British Isles. Taking the Irish, Scottish, and English specimens of the *Naias*, they seem to accord with the American plant, and, so far as I have seen, specimens show no approach to the var. *microcarpa* Nilsson from Lake Ruigsjön in Scania, Sweden. Of this I possess original specimens from Herr Nilsson. This is a smaller plant than ours—4-10 cm. to 1-50 dm. high, with fruit only 2—2.25 mm. long. In Europe *Naias* occurs in Finland; Russia, in Lithuania, Olentz, and Borussia; Germany, in Pomerania, Marchin! (Pasteiner-See, &c.); Sweden, Upland! (formerly) and Scania!; N. America, from Labrador to New England States across to Oregon. This shows what may be expected when our English Lakes are investigated as the Scottish ones have been by Mr W. West."—A. BENNETT.

*Hydrilla verticillata* Casp. Esthwaite Water, v.-c. 69b, July 31, 1914. New to Britain. Grows in slightly coloured water at about 8 feet, and is invisible from the surface. Associated with it are *Naias flexilis*, *Pot. pusillus*, *Pot. Sturrockii*, and *Callitriche autumnalis*—the first named being dominant. *Elodea canadensis*, though abundant elsewhere in the lake, is not included in this interesting linear-leaved association. The plant is uniformly pale green in colour, extremely slender and brittle. The branching is mainly below, the long resultant erect shoots being nearly simple. The internodes are  $\frac{5}{8}$  inch to  $\frac{3}{4}$  inch (rarely 1 inch) in length and therefore much more distinct than those of *Elodea*. The leaves are most often in whorls of 5, but 3's and 4's are frequent, and near the base of the shoots whorls of 3 shorter and broader leaves are common. The leaves are narrowly linear, pellucid, usually just over  $\frac{1}{2}$  inch long, patent, acute, and minutely serrulate. The teeth are few, small, very acute, antrorse, extra marginal, more distant below, and best seen near the apex. I am informed that the leaves are more nearly entire than those of other European specimens of this species at Kew. I was unable to find flowers, but winter buds were prominently shown on specimens gathered later in the season.—W. H. PEARSALL. "*Hydrilla verticillata*, var. *gracilis*."—A. J. WILLMOTT *in lit.* "I refer it to var. *pomeranica* (Reichb.). See *B.E.C. Report* 1914, p. 22."—G. C. DRUCE.

*Eriocaulon septangulare* With. Craigga More, Galway W., August 14, 1913.—W. C. BARTON.

*Eleocharis uniglumis* Schultes. Slacks in dunes, Freshfield, S. Lanes, v.-c. 59, July 19, 1914.—W. G. TRAVIS.

*Eleocharis multicaulis* Sm. Name confirmed by Arthur Bennett. [Ref. No. 227.] Marsh 30 feet above sea level, Valley Burn, Rackwick, Hoy, Orkney, June 23 and September 9, 1914. Native. Common in several small marshes.—H. H. JOHNSTON.

*Scirpis filiformis* Savi, var. *monostachys* Hook. Freshfield, S. Lanes, v.-c. 59, July 17, 1914.—W. G. TRAVIS. Also sent from damp ground, sea cliffs, Milford-on-Sea, S. Hants, v.-c. 11, August 1914.—J. COMBER. "Yes."—E. S. MARSHALL.

*Scirpus fluitans* L. Near Yarnton, Oxon, June 1914. Sent in order to put on record a curious instance of plant occurrence. *S. fluitans* is one of our rarest species, being only known from two localities, neither of which has recently yielded it. The place where I found it last year was well known to me in the eighties, as it was then a shallow piece of water, rich in *Charas*. Since 1900 the water level of one area has sunk, and vegetation—*Carex*, *Sparganium*, *Iris*, &c., have asserted themselves. In the dry period when I visited it there was but little water, but the surface of the wet ground was covered with masses of this *Scirpus*, which must have been brought, probably by aquatic birds, and finding a congenial home, with little competition, was thus enabled to make in a few years this remarkable increase.—G. C. DRUCE.

*Eriophorum angustifolium* Roth, var. *alpinum* Gaudin = var. *minus* Koch = *E. gracile* Smith, non Roth. [Ref. No. 3943]. Plentiful in bogs, north-east corrie of Ben Chalum, Mid Perth, v.-c. 88, from 2300 to 2500 feet, July 20, 1914.—E. S. MARSHALL. "Yes, and as usual with Mr Marshall's gatherings beautifully preserved; strictly speaking I suppose the spikes should be pedunculate."—G. C. DRUCE.

*Carex vesicaria* L. Marshy places, Naseby reservoir, Northants, v.-c. 32, June 20, 1914. Spikelets appear to be much more closely sessile than usual.—G. CHESTER. "Spikelets appear more closely sessile than usual. Yes, this is so, but it varies greatly in this, the other extreme (f. *pendulina*) has the lowest spikes drooping with stalks three inches long."—A. BENNETT. "Very fine. In Scotland and Ireland it usually has similar short, stout, sub-sessile female spikelets."—E. S. MARSHALL.

× *Carex csomadensis* Simonkai in *Evum. Pl. Transs.* 556, 1887, = *C. riparia* × *vesicaria*. [Ref. No. 4919]. Marshy meadow, Grendon Underwood, Bucks, July 1911. In considerable abundance.

Some specimens shaded off to *vesicaria*; others approached *riparia*. It is evidently a rare hybrid.—G. C. DRUCE.

*Carex strigosa* Huds. Penn Wood, Bucks, June 1904.—G. C. DRUCE.

*Carex capillaris* L. On the Sugar-loaf Limestone of Cronkley Fell, v.-c. 65, altitude 2000 feet, June 9, 1914. Growing in close association with *Thalictrum alpinum* L., *Helianthemum canum* Baumg., var. *vineale* (Pers.) and *Viola Riviniana* Reichb., forma *minor* Murb.—J. CRYER.

*Carex binervis* Sm. Near the Duddon Estuary, 1912. Coll. D. LUMB, ex G. C. DRUCE.

*Carex distans* L. (Non *C. distans* v. *maritima* auct. = *C. neglecta* Degl.). Wet meadows in Parish of Tredington, Worcestershire (an inland station), July 1903. See *Report* 1903, p. 28.—Coll. F. TOWNSEND; comm. G. C. DRUCE.

*Carex fulva* Host. [Ref. No. 641.] Chippenham Fen, Cambridge, v.-c. 29, June 14, 1914.—G. C. BROWN.

*Carex Oederi* Retz., var. *elatior* Anderson *Cyp. Scand.* 25, 1849. Wicken Fen, Cambridgeshire, July 1904. L. H. Bailey's unfortunate attempt to displace *C. Oederi* Retz., and use it for a form of *C. flava* caused much confusion among British botanists, the effects of which have not yet ceased. The fact is *C. Oederi* has almost as close relation with *extensa* as it has with *flava*, and is a good species distinct from both. The more usual state is a very small plant especially fond of the gravelly margins of pools and lochs, but in our calcareous fen areas this robust plant occurs.—G. C. DRUCE.

*Carex polygama* Schkuhr (*C. Buxbaumii* Wahl.). Arisaig, Westernness, July 1903. Very local. The only known locality, since it appears to have become extinct at Harbour Island, Lough Neagh, where I vainly sought for it on two occasions in 1903.—G. C. DRUCE.

*Carex salina* Wahl. Wick, Caithness, July 1907.—G. C. DRUCE.

*Carex gracilis* Curt. [Ref. No. 640]. Chippenham Fen, Cambridge, v.-c. 29, June 14, 1914. One large tuft—specimen rather gone over, but characteristic.—G. C. BROWN. "A slender form. I did not see this on the Fen when with the late Mr Fryer, but we saw good *C. stricta* Good. *C. gracilis* is a local species in Cambridgeshire."—A. BENNETT. "Yes."—E. S. MARSHALL. "My specimen is an im-

perfect one, but from the early falling of the fruits (June 14) and the amount of filamentous material on the leaf sheaths—I would suggest comparing it with *C. elata*.”—G. C. DRUCE.

*Carex gracilis* Curt., var. *gracilescens* Almq.? Naseby reservoir, Northants, v.-c. 32, June 20, 1914. Differs much from type, but hardly sure if it can come under *gracilescens*.—G. CHESTER. “Agrees very closely with the Cambridgeshire specimens so named for me by Dr Almquist.”—A. BENNETT. “I believe so.”—E. S. MARSHALL. “Yes, a characteristic plant of the reservoir and a frequent form in Northamptonshire, where Mr Chester has been doing excellent work.”—G. C. DRUCE.

*Carex muricata* L. Meadow, Malvern Wells, Worcester, v.-c. 37, June 26, 1914.—Coll. R. F. TOWNDROW; comm. S. H. BICKHAM. “I put this under *C. Pairaei* F. Schultz, of which I have seen an authentic specimen at the British Museum. I have recently gathered it in Surrey, Berks, and Middlesex. It appears to be a plant of somewhat moister situations than *C. contigua* Hoppe (*C. muricata* auct.). It seems to be widely distributed in Britain, for I have seen specimens from at least seventeen English and Scotch, and one Irish county.”—A. B. JACKSON. “Is, I suppose, the segregate plant apart from *Pairaei*.”—A. BENNETT. “Yes, the true Linnean plant = *C. Pairaei* F. Schultz.”—E. S. MARSHALL.

*Carex contigua* Hoppe × *divulsa* Good.? [Ref. No. 4060.] Growing with the supposed parents on a grassy roadside near West Monkton, S. Somerset, v.-c. 5, June 20, 1914. Intermediate in characters. More advanced material could not be obtained, as all the herbage had been mown a week later. I believe that the suggested origin is right; if so, it appears to be a new hybrid for Britain.—E. S. MARSHALL. “I do not feel I can pass an opinion on these hybrids; the finder is so much better prepared to do so, seeing them *in situ*.”—A. BENNETT.

*Carex divulsa* × *vulpina*. Ditchside, Bransford, Worcester, v.-c. 37, June 15 and 22, 1914. The specimens are from the same locality as those sent last year. See *Report* 1913, p. 506.—Coll. R. F. TOWNDROW; comm. S. H. BICKHAM. “If so, it would seem that *divulsa* is the dominant plant in it.”—A. BENNETT. “I am by no means convinced that this is a hybrid. Does it ripen its fruit?”—G. C. DRUCE. Also sent from roadside ditch, Leigh Sinton, Worcester, v.-c. 37, July 28, 1914. These specimens are from a new locality to those of this plant sent last year.—Coll. R. F. TOWNDROW; comm. S. H. BICKHAM. “This was sent to me fresh; a good intermediate.”—E. S. MARSHALL. “Mihi valde dubia forsan mera forma (abnormis) *C. divulsa*.”—A. THELLUNG.

*Carex paniculata* L. [Ref. No. 75.] Marsh by river, Mundford, v.-c. 28, June 18, 1914.—F. ROBINSON.

*Carex paradoxa* Willd. [Ref. No. 72.] Shady marsh land, Hockham, v.-c. 28, July 11, 1914.—F. ROBINSON. "Yes, characteristic specimens. It was growing with *Deyeuxia neglecta* Kunth = *Calamagrostis stricta* Nutt. A new record for Norfolk, of a rare British species."—A. BENNETT. "Correct; fruit fully ripe."—E. S. MARSHALL. "Yes, not given in *Top. Bot.* for West Norfolk."—G. C. DRUCE.

*Carex chordorrhiza* Ehrh. Marsh near the Tubeg of Mudal, Sutherland, July 1907.—G. C. DRUCE.

*Spartina alterniflora* Lois. Hill Head, near Titchfield, Hampshire, September 13, 1914. These specimens came from its most easterly recorded station on Southampton Water, where it is apparently having a hard struggle for existence with *S. Townsendi*. Dr F. H. Arnold's Sussex station, further east, "Thorney, not far from Pilsey, September 18, 1900" (*Suss. Fl.*, ed. 2, 1907, p. 124) has never, I believe, been confirmed. Mr Standen and I failed to see it there this year. Most of the dried specimens of *S. alterniflora* that I have seen appear to have a narrower leaf than in *S. Townsendi*, but this, I think, is due to their becoming more decidedly involute than those of *Townsendi* when being dried. Although some care was taken to try and avoid this, members will see I have not been wholly successful.—C. E. SALMON. "Right. If *S. Townsendi* is really a hybrid between this and *S. stricta*, it is an exception to the general rule, being so much stouter and brighter green than either."—E. S. MARSHALL.

*Phalaris minor* Retz. Cultivated ground, Paradis, Guernsey, August 2, 1914.—W. C. BARTON.

*Phalaris canariensis* L. Waste ground, Bumpas Lane, Zealand Road, Chester, v.-c. 58, September 28, 1914.—C. WATERFALL.

*Alopecurus bulbosus* Gouan. [Ref. No. 35.] Thick marsh near sea, by Burgh Castle, Yarmouth, v.-c. 27, May 11, 1914.—F. ROBINSON. "Correct."—E. S. MARSHALL.

*Agrostis alba* L., var *stolonifera* L. Side of ditch, Keyhaven, S. Hants, v.-c. 11, August 1914.—J. COMBER.—"Very good."—E. S. MARSHALL. "Var. *prorepens* Koch, the Linnean *stolonifera* is doubtful and may be = *verticillata* Vill."—G. C. DRUCE.

*Calamagrostis canescens* Druce (*lanceolata* Roth). Damp wood, Thompson, v.-c. 28, July 4, 1914.—F. ROBINSON. Also from Mow Fen,

Shouldham, W. Norfolk, v.-c. 28, June 24, 1914.—J. E. LITTLE. "Yes (*C. canescens* Druce)."—E. S. MARSHALL.

*Deyouxia neglecta* Kunth. [Ref. No. 7541.] Near Loch Watton, Caithness, July 1907.—G. C. DRUCE. Also from [Ref. No. 71] shady marshland, Shropham Hundreds, Norfolk, v.-c. 28, July 11, 1914.—F. ROBINSON. Also from [Ref. No. 37] Loch Scarmclett, Caithness, July 1907. These are typical plants which are locally plentiful there.—G. C. DRUCE.

*Deyouxia neglecta* Kunth, var. *scotica*. [Ref. No. 36.] Loch Watton, Caithness, July 1907. This has more acuminate glumes and is the plant which was recorded as *strigosa* by Mr Arthur Bennett in *Journ. Bot.* 1885, p. 253. I went to visit Dick's locality for the plant which was called *lapponica* in Smiles' *Life of Robert Dick*, and found only this form growing there. Afterwards I saw it near Loch Scarmclett. It really approaches *strigosa* in appearance, but does not agree with it in the length of the callus hairs, which are of the length of the floret in *strigosa*, which has also a broader and laxer panicle. *D. strigosa*, Prof. Hackel thinks, is probably a hybrid of *epigeios* and *neglecta*, and he is quite confident in rejecting these as *strigosa*. It is sufficiently distinct from normal *neglecta* to warrant a varietal name, var. *scotica*, characterised as "Panicles larger and more diffuse than type, glumes longer, and more longly acuminate.—G. C. DRUCE.

*Gastridium ventricosum* Sch. & Th. Copyhold, Sussex, September 1902. Once again the name must be changed in order to comply with the *Actes*. The trivial *lendigerum* dates from *Sp. Pl.*, ed. ii., 1762 or 1763, but prior to that it was published as *Agrostis ventricosa* by Gouan, and must stand as *G. ventricosum* (Gouan) Thellung.—G. C. DRUCE.

*Apera Spica-venti* Beauv. By River Orwell, Freston, v.-c. 25, July 13, 1913.—G. C. BROWN.

*Apera interrupta* Beauv. [Ref. No. 79.] Oatfield, Tottington, v.-c. 28, July 31, 1914.—F. ROBINSON. "*Apera Spica-venti* Beauv.—J. CRYER and W. G. TRAVIS. "This is *A. Spica-venti* Beauv., var. *purpurea* Rouy, distinguished not only by its purplish tint, but by the outer and inner glumes being longer and more acuminate. Compare Mr G. C. Brown's *Apera Spica-venti* Beauv., from Freston, v.-c. 25, which has paler, shorter glumes (especially the inner), and is the sub-var. *virescens* Rouy."—C. E. BRITTON. "Surely this is *A. Spica-venti*. Panicle large, broad, brownish, not interrupted."—E. S. MARSHALL. Also from sandy wayside, near Barnack, Northants, v.-c. 32, July 5, 1914.—G. CHESTER. "I am afraid only adventitious in Northants, and of course so in South Lanes."—G. C. DRUCE. Also

from sandy roadside, Freshfield, v.-c. 59, July 5, 1914.—J. A. WHELDON and W. G. TRAVIS. Also from sandy land, Cockley Cley, Swaffham, W. Norfolk, v.-c. 28, June 23, 1914.—J. E. LITTLE. "Fine and characteristic."—E. S. MARSHALL.

*Corynephorus canescens* Beauv. The Denes, Great Yarmouth, E. Norfolk, v.-c. 27, 1913.—Coll. Dr F. LONG; comm. G. C. BROWN. Also from North Denes, Great Yarmouth, v.-c. 27, August 13, 1914.—F. ROBINSON. "Yes, the var. *maritimus* Godron *Fl. Fr.* iii, p. 502. The common British plant."—G. C. DRUCE.

*Aira caryophyllea* L., *forma*. Petit Bo Cliffs, Guernsey, April 1907. On ground where gorse had been burned the previous year. Growing in prostrate tufts, much resembling *Aira praecox*.—G. C. DRUCE. "Only in bud; too young to afford grounds for a valid opinion."—E. S. MARSHALL.

*Deschampsia flexuosa* L., b. *montana* Hook. [Ref. No. 67]. Heath land amongst *Calluna*, Thompson, v.-c. 28, July 4, 1914.—F. ROBINSON. "No; only type. Var. *montana* is quite alpine."—E. S. MARSHALL. "A heath form. In the true *montana* the panicle is closed in fruit and the glumes are rich purple."—G. C. DRUCE.

*Arrhenatherum elatius* M. and K., var. *biaristatum* Druce. Near Lydd, Kent, July 1914, and the first species to grow on the shingle beds—that is, it approaches more closely to the sea than any other plant.—G. C. DRUCE.

*Koeleria gracilis* Pers. [Ref. No. 77]. Newmarket Heath, v.-c. 29, June 8, 1913. Is this type? It seems to be intermediate between *gracilis* and *britannica* in some respects.—W. C. BARTON. "Material too meagre; but I think it rightly named."—E. S. MARSHALL.

*Koeleria*—? [Ref. No. 112]. Sandy coast, Deal, East Kent, v.-c. 15, July 29, 1913.—W. C. BARTON. "A small form of *K. albescens* DC.; new for Kent, I think."—E. S. MARSHALL. "I wish the lower leaves on my specimen had been better shown. The plant suggests *K. albescens* DC. In that species the radical leaves and the leaves of the off-shoots are not flat as in *gracilis*, but enrolled-subulate. On the whole, I think it is very likely to be *albescens*; if so, a new county record."—G. C. DRUCE.

*Koeleria vallesiana* Asch. and Graebn. Uphill and Brean Down, N. Somerset, July 8, 1913. This grass seems to have recovered ground; there was a great quantity of it at Uphill, and a limited quantity on Brean Down, but only a small proportion was in flower.—H. J. RIDDELSDELL.

*Poa bulbosa* L. [Ref. No. 36]. Loose sand by sea, South Denes, Yarmouth, v.-c. 27, May 9, 1914.—F. ROBINSON.

*Glyceria festuciformis* Heyn., var. *hibernica*. Strangford Lough, Co. Down, August 1909. In considerable quantity. Not quite the continental species, differing in several points, especially in the less acuminate glumes.—G. C. DRUCE.

*Festuca rigida* Kunth, *forma* or *var.*? [Ref. No. 120]. Quarry Wood, Berks, v.-c. 22, June 22, 1913. A pretty form, growing on a bank under beech trees. Is it a usual shade form, or more than that? —W. C. BARTON. "Very curious; probably a starved state of dry woodlands."—E. S. MARSHALL.

*Festuca dumetorum* L., *forma planifolia* Hackel, *in lit.* comb. nov. [Ref. No. 4727.] Skegness, Lincoln, July 1911. Growing with the type. The occurrence of this flat-leaved form necessitates an alteration in description given in my edition of *Hayward's Pocket Book*, in which the leaves (of the type *dumetorum*) are described as "filiform"; it should read "filiform or flat, acute." Although there treated as a variety of *F. rubra* (for the sake of uniformity) *dumetorum* is a good species, and is so kept by Rouy *Fl. Fr.* xiv., p. 202, who, however, puts under it as a race, *F. arenaria* Osb., which we place under *F. rubra*.—G. C. DRUCE.

*Festuca ambigua* Le Gall? Blown sand, Pagharn, W. Sussex, v.-c. 13, June 15, 1914.—J. E. LITTLE. "*Festuca Myuros* L."—J. CRYER. "Certainly; just like the blown sand Deal to Sandwich plants."—E. S. MARSHALL. "Is *F. ambigua* Le Gall., the *F. Danthonii* A. & G., var. *ambigua* mihi, of my *List*. I found the type plentifully in Greece in 1914."—G. C. DRUCE.

*Festuca Myuros* L. [Ref. No. 644.] Berechurch, N. Essex, v.-c. 19, May 31, 1914.—G. C. BROWN. "Not *F. Myuros* L., but *F. bromoides* L. (*F. sciuroides* Roth)."—C. E. BRITTON and E. S. MARSHALL. "Not *Myuros*, but *F. bromoides* L., which is, oddly enough, only given sub-specific rank by Rouy. The fruits of the two plants are quite distinct."—G. C. DRUCE.

*Bromus maximus* Desf. [Ref. No. 42.] Sandhills by sea, North Denes, Great Yarmouth, v.-c. 27, May 10, 1914.—F. ROBINSON. "Hackel, and Rouy (*Fl. Fr.*) use the name *B. villosus* Forsk. for this, but *B. rigens* L. is older, but perhaps less well defined."—G. C. DRUCE.

*Bromus tectorum* L. [Ref. No. 52.] Roadside, light soil, three miles out of Thetford on London Road, West Suffolk, June 4, 1914.—F. ROBINSON. "Yes."—G. C. DRUCE.

*Bromus erectus* Huds., var. *glabrifolius* Borbas. Railway side, Seascale, v.-c. 70, June 5, 1914. I am not sure that the varietal name is correctly applied to these plants.—D. LUMB. "Yes."—G. C. DRUCE.

*Bromus unioloides* H. B. K. Waste ground, Hythe Quay, Colchester, v.-c. 19, June 2, 1913.—G. C. BROWN. "Yes."—G. C. DRUCE.

*Bromus secalinus* L., var. *hirtus* A. & G. Iver, Bucks, July 1903.—G. C. DRUCE.

*Bromus* sp. Mud wall top, Wigginton village, Oxon, June 23, 1914. Closely related to *B. racemosus*, but the compact panicle apparently separates it from that species.—H. J. RIDDELSDELL. "A curious, capitate form of *B. hordeaceus* L.; less hairy than the type, thus approaching var. *leptostachys (glabratus)*."—E. S. MARSHALL. "*B. hordeaceus* L., forma *congestus*."—G. C. DRUCE.

*Bromus arvensis* L. In sainfoin, Purwell, Hitchin, Herts, v.-c. 20, July 16, 1914. In the early state the palea is about 5 mm. shorter than the flowering glume. Later, in fruit, it equals or slightly exceeds it.—J. E. LITTLE. Also [Ref. No. 89] from farm roadway, Holme Hall, v.-c. 28, August 27, 1914.—F. ROBINSON. "Yes; Rouy puts it in the genus *Serrafalcus*, which also contains *secalinus*, *commutatus*, *racemosus*, *hordeaceus (mollis)* and my *Bromus interruptus = Serrafalcus interruptus mihi = S. pseudo-velutinus* Groves."—G. C. DRUCE.

*Agropyron pungens* R. & S. Shoreham, Sussex, June 1891.—G. C. DRUCE.

*Agropyron pungens* R. & S., forma *crisatum*. Teste Hackel. [Ref. No. 50]. Albecq and Vazon Bay, Guernsey, August 16, 1912. This form occurs in some quantity on the shingle inside the sea wall between Albecq and Vazon Bay. The plants seem to bear the same relation to the type as *Lolium perenne*, var. *crisatum* Doell, to type. Of this I sent some specimens last year, and have since noted some normal plants of *Lolium perenne* developing this *crisatum* form of spike, apparently as a result of damage through being trodden upon. However, these Guernsey plants are not in any sense damaged, though they very probably are starved. The type was plentiful thirty yards further inland.—W. C. BARTON. "Yes, Prof. Hackel so named similar specimens which I collected in the same locality in 1906."—G. C. DRUCE.

*Agropyron junceum* × *repens* Beauv. Keyhaven, S. Hants, v.-c. 11, August 1914.—J. COMBER. "Not in the least like any of my

specimens of this hybrid, mostly certified by Hackel. Why not *A. pungens*?—E. S. MARSHALL. "I think only luxuriant *pungens*."—G. C. DRUCE.

*Hordeum violaceum* Boissier. Selkirk, v.-c. 79, October 1913. Alien, Asia Minor. Det. A. Hackel. In this later gathering of 25th October, the beautiful blue shade is not so apparent as in my August specimens, owing possibly to the want of sunshine.—I. M. HAYWARD. "Yes, a beautiful grass, which, through Miss Hayward's generosity, was figured in our last *Report*."—G. C. DRUCE.

*Athyrium Filix-femina* Roth, var. *incisum* Newm. [Ref. No. 4804]. Four Slips Copse, Northants, August 1903.—G. C. DRUCE. "Yes."—F. W. STANSFIELD.

*Polystichum angulare* Presl, ? var. *Braunii* Spenner. Hedgebank near Newquay, Cardiganshire, May 14, 1914.—J. W. WHITE. "The more slender of these fronds is typical *angulare*. The denser one has some resemblance to *Braunii*, but in my opinion is a setose form of *angulare* going very slightly in the direction of *aculeatum*. The true *Braunii* of the U.S. America is very easily distinguished in the young growing (spring) state by both surfaces as well as rachis being thickly clothed with fine linear scales which are shed when the frond reaches maturity. I shall be pleased to send Mr White a fresh frond of true *Braunii* later in the season if he cares for it. I am sceptical of *Braunii* as a British fern. It seems to me to have affinities with *P. setosum*, although quite distinct from it."—F. W. STANSFIELD. "I have never seen this; but Mr White's plant differs greatly from ordinary *P. angulare*, which abounds in my parish, and agrees fairly well with the descriptions of Koch and Rouy. *Aspidium Braunii* Spenner is retained as a full species by Nyman, under the earlier, but preoccupied name *A. angulare* Kit.; Rouy makes it a sub-species."—E. S. MARSHALL.

*Dryopteris cristata* A. Gray. Scoulton, Norfolk, August 1904.—G. C. DRUCE. "Yes."—F. W. STANSFIELD.

*Lastrea glandulosa* Newm. Ankerbury Bog, v.-c. 34, June 2, 1913. Stipes usually thickly glandular; fronds glandular beneath but not very thickly. The scales are not as dark as those of *dilatata*; the fronds may be described as broadly lanceolate: at any rate they are not those of true *dilatata*. In a very wet piece of wood below a bog on Ankerbury Hill, Lydbrook, W. Glos. Whether this is the place first made famous by Mr Purchas's discoveries or not, I am uncertain; the houses now come within a hundred yards of it, and the bog has shrunk. A number of circumstances make me doubt whether the name is correct; and I have never seen an authenticated

specimen of Newman's *glandulosa*.—H. J. RIDDELSDELL.—“*L. glandulosa* Newm. is described by Newman as having concolorous scales, while those of this specimen are maculated. *Glandulosa* and *collina* Newm. are both forms intermediate between *dilatata* (*aristata*) and *spinulosa*. There are many such intermediate forms, and they are by no means well defined. I should call this specimen *L. collina* Newm., as having somewhat of the habit of *spinulosa* with the bicolorous scales of *dilatata*.”—F. W. STANSFIELD.

*Dryopteris rigida* Underw. Hutton Roof, Westmoreland, August 1909.—G. C. DRUCE. “I agree. It was very abundant forty years ago in the station.”—F. W. STANSFIELD.

*Hymenophyllum tunbridgense* Sm. Rocky wood in W. Glos., June 1, 1914. Coll. H. H. KNIGHT. A few scraps just to establish the record. I believe I recorded it somewhere in 1911, when Mr Knight first showed me the plant. He tells me now that the fern is in fair quantity on rock faces, in very dark parts of the wood. Some of the sheets bear fruit.—H. J. RIDDELSDELL. “Yes.”—F. W. STANSFIELD.

*Nitella opaca* Agardh. Looe Pool, Cornwall, July 1904.—G. C. DRUCE.

*Nitella hyalina* Agardh. Carminow Creek, Looe Pool, Helston, W. Cornwall, August 28, 1914. I think this is the right plant, though they are smaller than other specimens I have.—Coll. E. THURSTON; comm. C. C. VIGURS.

*Tolypella glomerata* Leonh. Marcham, Berks, June 1891.—G. C. DRUCE.

---

#### CORRECTIONS IN DISTRIBUTOR'S REPORT FOR 1913.

Page 444. *Ranunculus Flammula* L. In second line, for “Kirbistry” read “Kirbister.”

Page 471. *Hedera Helix* L., var. *borealis* Druce. In fifth line omit “Lieutenant,” and in sixth line, for “Bethsdale” read “Berriedale.”

