

## NOMENCLATURE AND CORRECTIONS TO BRITISH PLANT LIST

## VII

By E. F. WARBURG.

This feature was in course of preparation by the late Mr A. J. Wilmott at the time of his death. At present, only his rough notes are available. The following list is therefore confined to additions and changes made necessary by material published in *Watsonia* during the past year, including some name changes made in Plant Records which are explained below. References cited only by page numbers refer to *Watsonia*, 1.

- 44           **EROPHILA.**  
5   *conferta* Wilmott. See paper by Wilmott, pp. 137-138.
- 170           **CORONILLA**  
4   *glauca* L. See Plant Notes.
- 185           **RUBUS.** See paper by Watson, pp. 71-83, unless another reference is given.
- 9(2)   **Briggsianus** (Rog.) W. Wats.—1946, *J. Ecol.*, **33**, 337.  
Vice 9b.
- 18(4)   **latiarcuatus** W. Wats.
- [19   **argenteus** Weihe & Nees. Delete.]
- 41(2)   **sciocharis** (Sudre) W. Wats.—1946, *J. Ecol.*, **33**, 339; *R. gratus* ssp. *R. sciocharis* Sudre, 1907, *Bat. Eur.*, 68.  
Vice *R. sciaphilus* Lange, 1883, non Lef. & Muell., 1859 (see 1938, *Rep. Bot. Soc. and Exch. Club*, **11**, 445) and 41b. (*B.P.L.*).
- 53(3)   **Libertianus** Weihe.
- 56(3)   **albionis** W. Wats. Vice 56 b.
- 61   **bracteosus** Weihe—in Lej. & Court., 1831, *Comp. Fl. Belg.*, **2**, 162; *R. orthocladus* A. Ley, 1896. See Watson, 1946, *J. Ecol.*, **33**, 340.
- 65(2)   **macroacanthos** Weihe & Nees.
- 67(4)   **hypomalacus** Focke.
- 67(5)   **silesiacus** Weihe.
- 95(2)   **Watsonii** Mills. See paper by Mills, pp. 135-136.
- 98(3)   **melanoxylon** Muell. & Wirtg.
- 98(4)   **Reichenbachii** Koehl. ex Weihe.
- 102   **breconensis** W. Wats. *R. Lejeunei* Rogers, *Handbook*, 70, non Weihe.
- 102(3)   **Lejeunei** Weihe.
- 108(3)   **Menkei** Weihe.

- 110(5) **fusciformis** Sudre.  
 111(2) **adornatiformis** (Sudre) Bouv.  
 120(2) **foliosus** Weihe.  
 122(3) **scabripes** Genev. Vice 122c.  
 130(2) **humifusus** Weihe.  
 137 **angustifrons** Sudre. Delete brackets.  
 [139 **serpens**. Delete.]  
 140(2) **Guentheri** Weihe.  
 147(2) **pygmaeus** Weihe.  
 149(3) **purpureicaulis** W. Wats. Vice 149h. See paper by  
 Watson, pp. 289-290.  
 149(4) **Babingtonianus** W. Wats. See paper by Watson, pp.  
 289-290.
- 190(2) APHANES** L. See paper by Walters, pp. 163-169.  
 1 **arvensis** L. Vice 190/19.  
 2 **microcarpa** (Boiss. & Reut.) Rothm.
- 299(1) PHUOPSIS** Benth. & Hook. f. See Polunin, 1942, *Rep.  
 Bot. Soc. and Exch. Club*, **12**, 356-358.  
 1 **stylosa** (Trin.) Benth. & Hook. f. ex Jacks. N.W. Persia.  
 Vice 299/3 *Crucianella stylosa* Trin. (see 1946, *Rep.  
 Bot. Soc. and Exch. Club*, **12**, 663).
- 354 GALINSOGA.**  
 2 **ciliata** (Raf.) Blake. Vice 2b. *G. quadriradiata* var. *his-  
 pida* (DC.) Thell. (see 1939, *Rep. Bot. Soc. and Exch.  
 Club*, **12**, 44, 94). See paper by Lousley, pp. 238-241.
- 411 LAPSANA.**  
 2 **intermedia** M. Bieb. See paper by Burt, pp. 234-237.
- 416 CREPIS.** See Review, pp. 195-199.  
 1 **mollis** (Jacq.) Asch.—1864, *Fl. Brandenb.*, 386; *Hiera-  
 cium molle* Jacq., 1774, *Fl. Austr.*, **2**, 12.  
 9 **foetida.**  
 A. **eufoetida** Beger ex Domin—1935, *Preslia*, **13-15**,  
 252. Vice subsp. *vulgaris* (Bisch.) Babc., 1938  
 (see 1939, *Rep. Bot. Soc. and Exch. Club*, **12**,  
 26) and a. *vulgaris* Bisch. (see 1934, *Rep. Bot.  
 Soc. and Exch. Club*, **10**, 485).  
 B. **rhoadifolia** (Bieb.) Schinz & Thell.—in Schinz &  
 Keller, 1914, *Fl. Schweiz*, ed. 3, **2** (*Krit. Fl.*),  
 361. Vice 3b. *hispida* Bisch. (see 1934, *Rep.  
 Bot. Soc. and Exch. Club*, **10**, 485).  
 C. **commutata** (Spreng.) Babc.—1938, *J. Bot.*, **76**, 207.  
 Vice 414/1.  
 13 **sancta** (L.) Babc.—1941, *Univ. Calif. Publ. Bot.*, **19**, 403;  
*Hieracium sanctum* L., 1756, *Cent. Plant.*, **2**, 30.  
 Vice 418/1.

- 14 *Zacintha* (L.) Babc.—1941, *Univ. Calif. Publ. Bot.*, **19**, 404; *Lapsana Zacintha* L., 1753, *Sp. Plant.*, 811. Vice 430(2)/1 *Zacyntha verrucosa* Gaertn. (see 1932, *Rep. Bot. and Exch. Club*, **10**, 27).
- 416(2)**     **AETHÉORRHIZA** Cass.—1827, *Dict. Sci. Nat.*, **48**, 425.  
 1 *bulbosa* (L.) Cass.—1827, *loc. cit.* Vice *Crepis bulbosa* (L.) Tausch (see 1929, *Rep. Bot. Soc. and Exch. Club*, **8**, 618, 880). See Review, p. 199, and Babcock & Stebbins, 1943, *Univ. Calif. Publ. Bot.*, **18**, 235-240.
- 558**             **MENTHA.**  
 3 *longifolia*  
     i. *horridula* Briquet. Vice 3×4=*Nouletiana*. See paper by Graham, p. 278.
- 566**             **SALVIA.**  
 [21 *lanceolata* Willd. (see 1939, *Rep. Bot. Soc. and Exch. Club*, **12**, 53). Delete ; =20 *reflexa* Hornem. (see 1929, *Rep. Bot. Soc. and Exch. Club*, **8**, 634). See Plant Notes.]
- 615**             **POLYGONUM.**  
 25 *cognatum*. See Plant Notes.  
     a. *alpestre* (C. A. Mey.) Meisn.  
     b. *ammanioides* (Jaub. & Spach) Meisn.
- 650**             **SALIX.** See paper by Rechinger, pp. 271-275.  
     6(2) *dasyclados* Wimm. Vice "6 [×] *acuminata* (Sm.)."  
     6(2) × 5 = *Taylori* Rech.  
     10 × 8 × 6  
     11(2) × 6  
     11(2) × 10 × 6 = *angusensis* Rech.
- 668**             **EPIPACTIS.**  
 3(5) *cambrensis* C. Thomas. See paper by Thomas, pp. 283-288.
- 683**             **CROCOSMIA** Planch.—1851-2, *Flores des Serres*, **7**, 161.  
 1 × *crocosiiflora* (Lemoine) N.E.Br.—1932, *Trans. R. Soc. S. Afr.*, **20**, 264 = *C. aurea* (Pappe) Planch. × *C. Pottsii* (Baker) N.E.Br.; *Montbretia crocosmiæflora* Lemoine [? *Cat.*, 1881] et ex Morren, 1881, *La Belgique Horticole*, **31**, 299, et apud André, 1882, *Rev. Hort.*, **54**, 124; *Tritonia crocosmiiflora* (Lemoine) Nichols., 1887, *Dict. Gard.*, **4**, 94. Transferred from *Tritonia* to *Crocoshmia* by N. E. Brown, *loc. cit.*—From A. J. Wilmott and H. K. Airy Shaw.

744

**CYPERUS.**

- 4 *Eragrostis* Lam., 1791, *Tabl. Encycl. Bot.*, **1**, 196; *C. declinatus* Moench, 1794, (*B.P.L.*); *C. vegetus* Willd., 1798, *Sp. Pl.*, 283 (*nomen abortivum*).—A. J. Wilmott.

844

**EQUISETUM.** See paper by Alston, pp. 149-153.

- 7(2) **Moorei** Newm.—1854, *Phytologist*, 19. Vice 7b. and 7(2) *occidentale* (Hy) Coste (see 1929, *Rep. Bot. Soc. and Exch. Club*, **9**, 42).
- 7(3) **ramosissimum** Desf.

876

**CHARA.**

- 16 **globularis** Thuill. (1799, *Fl. Env. Paris*, ed. 2, 472) emend. Zanev. (1940, *Blumea*, **4** (1), 191). Vice *C. fragilis* Desv. in Loisel., 1810, *Not. Fl. Fr.*, 137.

a. **capillacea** (Thuill.) Zanev.—1940, *loc. cit.*, 195; *C. capillacea* Thuill., 1799, *loc. cit.*, 474; *C. fragilis* Desv., 1810, *loc. cit.*, *sensu stricto*; *C. pulchella* Wallr. var. *capillacea* (Thuill.) Wallr., 1833, *Fl. Crypt. germ.*, **2**, 109.

b. **globularis** (Thuill.) Wilmott, comb. nov.; *C. globularis* Thuill., 1799, *loc. cit.*, *sensu stricto*; *C. pulchella* Wallr. var. *globularis* (Thuill.) Wallr., 1833, *loc. cit.*; *C. Hedwigii* Ag. ex Bruz., 1824, *Obs. Char.*, **7**, 21; *C. fragilis* Desv. var. *Hedwigii* (Ag. ex Bruz.) Kütz., 1843, *Phyc. Gen.*, 319.

Zaneveld (1940, *loc. cit.*) appears to have been the first to combine Thuillier's two species under one of his names.—From G. O. Allen and A. J. Wilmott.